



US009359628B2

(12) **United States Patent**
Contreras et al.

(10) **Patent No.:** US 9,359,628 B2
(45) **Date of Patent:** *Jun. 7, 2016

(54) **PROTEIN GLYCOSYLATION
MODIFICATION IN METHYLOTROPHIC
YEAST**

(71) Applicants: **VIB, vzw**, Zwijnaarde (BE); **Research Corporation Technologies, Inc.**, Tucson, AZ (US); **Universiteit Gent**, Ghent (BE)

(72) Inventors: **Roland Contreras**, Merelbeke (BE); **Nico L.M. Callewaert**, Lichtervelde (BE); **Steven C.J. Geysens**, Kruishoutem (BE)

(73) Assignees: **VIB, VZW**, Zwijnaarde (BE); **Research Corporation Technologies, Inc.**, Tucson, AZ (US); **Universiteit Gent**, Ghent (BE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **14/152,329**

(22) Filed: **Jan. 10, 2014**

(65) **Prior Publication Data**

US 2014/0154738 A1 Jun. 5, 2014

Related U.S. Application Data

(63) Continuation of application No. 12/424,924, filed on Apr. 16, 2009, now Pat. No. 8,663,971, which is a continuation of application No. 10/672,484, filed on Sep. 25, 2003, now Pat. No. 8,354,268, which is a continuation of application No. 09/896,594, filed on Jun. 29, 2001, now Pat. No. 6,803,225.

(60) Provisional application No. 60/215,676, filed on Jun. 30, 2000.

(51) **Int. Cl.**

C12N 1/14 (2006.01)
C12N 15/74 (2006.01)
C12P 21/04 (2006.01)
C12P 21/00 (2006.01)
C12N 15/81 (2006.01)
C12R 1/84 (2006.01)
C12N 9/24 (2006.01)
C07H 21/04 (2006.01)
A61K 38/00 (2006.01)

(52) **U.S. Cl.**

CPC **C12P 21/005** (2013.01); **C12N 9/2402** (2013.01); **C12N 9/2488** (2013.01); **C12N 15/81** (2013.01); **C12N 15/815** (2013.01);
C12R 1/84 (2013.01); **C12Y 302/0102** (2013.01); **C12Y 302/01024** (2013.01); **A61K 38/00** (2013.01); **Y02P 20/52** (2015.11)

(58) **Field of Classification Search**

CPC C12P 21/005; C12N 15/81; C12N 15/815;
C12R 1/84

USPC 435/254.23, 483, 71.1
See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,135,854 A	8/1992	MacKay et al.
5,705,616 A	1/1998	Lehle et al.
6,803,225 B2	10/2004	Contreras et al.
7,252,933 B2	8/2007	Contreras et al.
7,507,573 B2	3/2009	Contreras et al.
8,058,053 B2	11/2011	Contreras et al.
8,354,268 B2 *	1/2013	Contreras et al. 435/254.23
8,663,971 B2 *	3/2014	Contreras et al. 435/254.23
8,883,445 B2 *	11/2014	Contreras et al. 435/69.1
2002/0137134 A1	9/2002	Gerngross

FOREIGN PATENT DOCUMENTS

EP	0 314 096 A2	5/1989
EP	1 211 310 A1	6/2002
EP	1 297 172 B1	4/2003
JP	8-336387	12/1996
JP	9-261	1/1997
WO	91/05057	4/1991
WO	96/21038	7/1996
WO	02/00879 A2	1/2002
WO	2004/003205 A1	1/2004

OTHER PUBLICATIONS

Grinna, L.S. et al., "Size Distribution and General Structural Features of N-linked oligosaccharides from the methylotrophic yeast, *Pichia pastoris*" Yeast (Mar.-Apr. 1989) pp. 107-115, vol. 5, No. 2.

Jungmann, J. et al., "The *Saccharomyces cerevisiae* protein Mnn10p/Bed1p is a subunit of a Golgi mannosyltransferase complex" J Biol Chem. (Mar. 5, 1999) pp. 6579-6585, vol. 274, No. 10.

Lee, B.N. et al., "The MAPKKK Ste11 regulates vegetative growth through a kinase cascade of shared signaling components" PNAS (Oct. 26, 1999) pp. 12679-12684, vol. 96, No. 22.

Nakayama, K. et al., "Substrate specificity of α -1,6-mannosyltransferase that initiates N-linked mannose outer chain elongation in *Saccharomyces cerevisiae*" FEBS Letters (Aug. 4, 1997) pp. 547-550, vol. 412, No. 3.

Becker, B. et al., "The transmembrane domain of murine alpha-mannosidase IB is a major determinant of Golgi localization" European J. Cell Biol (Dec. 2000) pp. 79: 986-992 vol. 79, No. 12.

Choi, B.K. et al., "Use of combinatorial genetic libraries to humanize N-linked glycosylation in the yeast *Pichiapastoris*" PNAS (Apr. 29, 2003) pp. 5022-5027, vol. 100, No. 9.

Conzalez, D.S. et al., "The α -Mannosidases: Phylogeny and Adaptive Diversification" Mol Biol Evolution (Nov. 2000) pp. 292-300, vol. 17, No. 2.

(Continued)

Primary Examiner — Quang Nguyen

(74) **Attorney, Agent, or Firm — Scully, Scott, Murphy & Presser, P.C.**

(57)

ABSTRACT


The present invention provides genetically engineered strains of *Pichia* capable of producing proteins with reduced glycosylation. In particular, the genetically engineered strains of the present invention are capable of expressing either or both of an α -1,2-mannosidase and glucosidase II. The genetically engineered strains of the present invention can be further modified such that the OCH1 gene is disrupted. Methods of producing glycoproteins with reduced glycosylation using such genetically engineered stains of *Pichia* are also provided.

8 Claims, 23 Drawing Sheets

(56)

References Cited**OTHER PUBLICATIONS**

- Herscovics, A., "Structure and Function of Class I α 1,2-mannosidases Involved in Glycoprotein Synthesis and Endoplasmic Reticulum Quality Control" *Biochemie* (Aug. 2001) pp. 757-762, vol. 83, No. 8.
- Higgins, D.R. et al., Methods in Molecular Biology in *Pichia* Protocols, Higgins and Gregg (eds.) 1998, 10 pages.
- Nett, J.H. et al., "A combinatorial genetic library approach to target heterologous glycosylation enzymes to the endoplasmic reticulum or the Golgi apparatus of *Pichia pastoris*" *Yeast* (Jan. 2011) pp. 237-252, vol. 28.
- Schneikert, J. et al., "Characterization of a novel mouse recombinant processing α -mannosidase" *Glycobiology* (Apr. 1994) pp. 445-450, vol. 4, No. 4.
- Bretthauer, R.K. et al., "Glycosylation of *Pichia pastoris*-derived Proteins" *Biotechnol. Appl. Biochem.* (Dec. 1999) pp. 193-200, vol. 30.
- Ceregbino J.L. et al., "Heterologous Protein Expression in the Methylotrophic Yeast *Pichia Pastoris*" *FEMS Microbiology Reviews* 24:45-66 (2000).
- Herscovics A., "Processing Glycosidases of *Saccharomyces cerevisiae*" *Biochimica et Biophysica Acta* 1426:275-285 (1999).
- Kang H.A. et al., "Glycostylation of Human α 1-Antitrypsin in *Saccharomyces cerevisiae* and Methylotrophic Yeasts" *Yeast* 14:371-381 (1998).
- Tremblay L.O. et al., "Molecular Cloning, Chromosomal Mapping and Tissue-Specific Expression of a Novel Human α 1,2-Mannosidase Gene involved in N-Glycan Maturation" *Glycobiology* 8:585-595 (1998).
- Malissard M. et al., "The Yeast Expression System, for Recombinant Glycosyltransferases" *Glycoconjugate Journal* 16:125-139 (1999).
- Nagasu T. et al. "Isolation of New Temperature-Sensitive Mutants of *Saccharomyces cerevisiae* Deficient in Mannose Outer Chain Elongation", *Yeast* 8:535-547 (1992).
- Vervecken W. et al., "In Vivo Synthesis of Mammalian-Like, Hybrid-Type N-Glycans in *Pichia Pastoris*", *Applied and Environmental Microbiology* 70(5):2639-2646 (2004).
- Trimble R.B. et al., "Structure of Oligosaccharides on *Saccharomyces* SUC2 Invertase Secreted by the Methylotrophic Yeast *Pichia Pastoris*", *The Journal of Biological Chemistry* 266(34)22807-22817 (1991).
- Verostek M.F. et al., "Mannosyltransferase Activities in Membranes from Various Yeast Strains", *Glycobiology* 5 (7):671-681 (1995).
- Pelham H.R.B. et al., "Sorting of Soluble ER Proteins in Yeast", *The EMBO Journal* 7(6):1757-1762 (1988).
- Blandin G. et al., "Genomic Exploration of the Hemiascomycetous Yeasts: 13. *Pichia Angusta*", *FEBS Letter* 487:76-81 (2000).
- Kim M.W. et al., "Functional Characterization of the Hansenula Polymorpha HOC1, OCH1, and OCR1 Genes as Members of the Yeast OCH1 Mannosyltransferase Family Involved in Protein Glycosylation", *The Journal of Biological Chemistry* 281(10):6261-6272 (2006).
- Ramezani-Rad M. et al., "The Hansenula Polymorpha (strain CES4732) Genome Sequencing and Analysis", *FEMS Yeast Research* 4:207-215 (2003).
- Alani E. et al., "A Method for Gene Disruption that Allows Repeated Use of URA3 Selection in the Construction of Multiply Disrupted Yeast Strains", *Genetics* 116:541-545 (1987).
- Nakayama Ken-ichi et al., "OCH1 encodes a novel membrane bound mannosyltransferase: outer chain elongation of asparagine-linked oligosaccharides", *The EMBO Journal* 11(7): 2511-2519 (1992).
- Kniskern P. J. et al., "Characterization and evaluation of a recombinant hepatitis B vaccine expressed in yeast defective for N-linked Hyperglycosylation". *Vaccine* 12(11): 1021-1025 (1994).
- Lehle L. et al., "Glycoprotein biosynthesis in *Saccharomyces cerevisiae*: ngl29, an N-glycosylation mutant allelic to och1 having a defect in the initiation of outer chain formation", *FEBS Letters* 370: 41-45 (1995).
- Yoko-o T. et al., "Schizosaccharomyces pombe och1+ encodes α -1,6-mannosyltransferase that is involved in outer chain elongation of N-linked oligosaccharides", *FEBS Letters* 489: 75-80 (2001).
- Cregg J. M. et al., "High-Level Expression and Efficient Assembly of Hepatitis B Surface Antigen in the Methylotrophic Yeast, *Pichia Pastoris*", *Biotechnology* 5: 479-485 (1987).
- Lai A. et al., "Substrate specificities of recombinant murine Golgi α 1,2-mannosidases IA and 1B and comparison with endoplasmic reticulum and Golgi processing α 1,2-mannosidases", *Glycobiology* 8(10): 981-995 (1998).
- Tremblay L. O. et al., "Cloning and expression of a specific human α 1,2-mannosidase that trims Man9GlcNAc2 to Man8GlcNAc2 isomer B during N-glycan biosynthesis", *Glycobiology* 9(10): 1073-1078 (1999).
- Gonzalez D. S. et al., "Identification, Expression, and Characterization of a cDNA Encoding Human Endoplasmic Reticulum Mannosidase I, the Enzyme That Catalyzes the First Mannose Trimming Step in Mammalian Asn-linked Oligosaccharide Biosynthesis", *The Journal of Biological Chemistry* 274(30): 21375-21386 (1999).
- Callewaert N. et al., "Use of HDEL-tagged *Trichoderma reesei* mannoseyl oligosaccharide α 1,2- α -D-mannosidase for N-glycan engineering in *Pichia pastoris*", *FEBS Letters* 503: 173-178 (2001).
- Maras, M., et al. (2000) "Molecular Cloning and Enzymatic Characterization of a *Trichoderma Reesi* 1, 2- α -D-Mannosidase", *Journal of Biotechnology* 77 255-263.
- Kukuruzinska M. A., et al. (1987) "Protein Glycosylation in Yeast", *Ann. Rev. Biochem* 56: 915-944.
- Chiba, Y., et al. (1998) "Production of Human Compatible High Mannose-Type (Man5GlcNAc2) Sugar Chains in *Saccharomyces cerevisiae*", *The Journal of Biological Chemistry* 273 (41): 26298-26304.
- Maras, M., et al. (1999) In Vivo Synthesis of Complex N-Glycans by Expression of Human N-Acetylglucosaminyltransferase I in the Filamentous Fungus *Trichoderma Reesi*, *FEBS Letters* 452: 365-370.
- Nakanishi-Shindo, Y., et al. (1993) "Structure of the N-Linked Oligosaccharides That Show the Complete Loss of α 1,6-Polymannose Outer Chain from och1, och1 mnn1, and och1 mnn1 alg3 Mutants of *Saccharomyces cerevisiae*", *The Journal of Biological Chemistry* 268 (35): 26338-26345.
- Martinet, W., et al. (1998) "Modification of the Protein Glycosylation Pathway in the Methylotrophic Yeast *Pichia Pastoris*", *Biotechnology Letters* 20(12):1171-1177.
- Maras, M., et al. (1997) "In Vitro Conversion of the Carbohydrate Moiety of Fungal Glycoproteins to Mammalian-Type Oligosaccharides", *Eur. J. Biochem.* 249:701-7707.
- Laroy,W., et al. (2000) "Cloning of Trypanosoma cruzi tans-Sialidase and Expression in *Pichia pastoris*", *Protein Expression and Purification* 20: 389-393.
- Inoue et al. Molecular cloning and nucleotide sequence of the 1,2-alpha-D-mannosidase gene, msdS, from *Aspergillus saitoi* and expression of the gene in yeast cells. *Biochim. Biophys. Acta* 1253:141-145, 1995.
- Herscovics et al. Isolation of a mouse Golgi mannosidase cDNA, a member of gene family conserved from yeast to mammals. *J. Biol. Chem.* 269:9864-9871, 1994.
- Lal et al. Isolation and expression of murine and rabbit cDNAs encoding an alpha 1,2-mannosidase involved in the processing of asparagine-linked oligosaccharides. *J. Biol. Chem.* 269:9872-9881, 1995.
- Trombetta et al. Endoplasmic reticulum glucosidase II is composed of a catalytic subunit, conserved from yeast to mammals, and a tightly bound noncatalytic HDEL-containing subunit. *J. Biol. Chem.* 271:27509-27516, 1996.
- Ngo et al. Computational complexity, protein structure prediction, and the Levinthal Paradox. IN: The protein folding problem and tertiary structure prediction (Merz et al., Eds.), Birkhauser, Boston, 1994, pp. 491-495.
- Rudinger J. Characteristic of the amino acids as components of a peptide hormone sequence. In: Peptide hormones (Parsons, J.A., Ed.), University Park Press, Baltimore, 1976, pp. 1-7.

**FIG. 1**

L929 (cycloheximide+ TNF induced) mouse cDNA library,
Group 6, average insert length : 2000 bp.




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11

GlsII Pichia expression vector pGAPZ(A)glsII




FIG. 12

GlsII Pichia expression vector pAOX2Z(A)glsII




FIG. 13

GlsII Pichia expression vector pPICZ(A)glsII




FIG. 14

GlsII Pichia expression vector pYPT1Z(A)glsII




FIG. 15

GlsII Pichia expression vector pGAPADE1glsII




FIG. 16

GlsII Pichia expression vector pAOX2ADE1glsII




FIG. 17

GlsII Pichia expression vector pPICADE1glsII




FIG. 18

GlsII Pichia expression vector pYPT1ADE1glsII




FIG. 19

GlsII Pichia expression vector pGAPZ(A)glsIIHDEL




FIG. 20

GlsII Pichia expression vector pGAPADE1glsIIHDEL




FIG. 21

Glucosidase II assay on commercially available alpha-glucosidase




FIG. 22

Glucosidase II assay on heterologously expressed Pichia protein




FIG. 23

1

**PROTEIN GLYCOSYLATION
MODIFICATION IN METHYLOTROPHIC
YEAST**

**CROSS-REFERENCE TO RELATED
APPLICATION**

This application is a continuation of U.S. application Ser. No. 12/424,924, filed Apr. 16, 2009, which is a continuation of U.S. application Ser. No. 10/672,484, filed Sep. 25, 2003, now U.S. Pat. No. 8,354,268, which is a continuation of U.S. application Ser. No. 09/896,594, filed Jun. 29, 2001, now U.S. Pat. No. 6,803,225, which claims the benefit of U.S. Provisional Application Ser. No. 60/215,676, filed Jun. 30, 2000.

FIELD OF THE INVENTION

The present invention relates to methods and vectors useful for genetically modifying the glycosylation process in methy lotrophic yeast strains for the purpose of producing glycoproteins with reduced glycosylation. The present invention further relates to methy lotrophic yeast strains generated using the present methods and vectors, as well as glycoproteins produced from such genetically modified strains.

BACKGROUND OF THE INVENTION

The methy lotrophic yeasts including *Pichia pastoris* have been widely used for production of recombinant proteins of commercial or medical importance. However, production and medical applications of some therapeutic glycoproteins can be hampered by the differences in the protein-linked carbohydrate biosynthesis between these yeasts and the target organism such as a mammalian subject.

Protein N-glycosylation originates in the endoplasmic reticulum (ER), where an N-linked oligosaccharide (Glc₃Man₉GlcNAc₂) assembled on dolichol (a lipid carrier intermediate) is transferred to the appropriate Asn of a nascent protein. This is an event common to all eukaryotic N-linked glycoproteins. The three glucose residues and one specific α-1,2-linked mannose residue are removed by specific glucosidases and an α-1,2-mannosidase in the ER, resulting in the core oligosaccharide structure, Man₈GlcNAc₂. The protein with this core sugar structure is transported to the Golgi apparatus where the sugar moiety undergoes various modifications. There are significant differences in the modifications of the sugar chain in the Golgi apparatus between yeast and higher eukaryotes.

In mammalian cells, the modification of the sugar chain proceeds via 3 different pathways depending on the protein moiety to which it is added. That is, (1) the core sugar chain does not change; (2) the core sugar chain is changed by adding the N-acetylglucosamine-1-phosphate moiety (GlcNAc-1-P) in UDP-N-acetyl glucosamine (UDP-GlcNAc) to the 6-position of mannose in the core sugar chain, followed by removing the GlcNAc moiety to form an acidic sugar chain in the glycoprotein; or (3) the core sugar chain is first converted into Man₅GlcNAc₂ by removing 3 mannose residues with mannosidase I; Man₅GlcNAc₂ is further modified by adding GlcNAc and removing 2 more mannose residues, followed by sequentially adding GlcNAc, galactose (Gal), and N-acetylneurameric acid (also called sialic acid (NeuNAc)) to form various hybrid or complex sugar chains (R. Kornfeld and S. Kornfeld, *Ann. Rev. Biochem.* 54: 631-664, 1985; Chiba et al *J. Biol. Chem.* 273: 26298-26304, 1998).

2

In yeast, the modification of the sugar chain in the Golgi involves a series of additions of mannose residues by different mannosyltransferases ("outer chain" glycosylation). The structure of the outer chain glycosylation is specific to the organisms, typically with more than 50 mannose residues in *S. cerevisiae*, and most commonly with structures smaller than Man₁₄GlcNAc₂ in *Pichia pastoris*. This yeast-specific outer chain glycosylation of the high mannose type is also denoted hyperglycosylation.

Hyperglycosylation is often undesired since it leads to heterogeneity of a recombinant protein product in both carbohydrate composition and molecular weight, which may complicate the protein purification. The specific activity (units/weight) of hyperglycosylated enzymes may be lowered by the increased portion of carbohydrate. In addition, the outer chain glycosylation is strongly immunogenic which is undesirable in a therapeutic application. Moreover, the large outer chain sugar can mask the immunogenic determinants of a therapeutic protein. For example, the influenza neuraminidase (NA) expressed in *P. pastoris* is glycosylated with N-glycans containing up to 30-40 mannose residues. The hyperglycosylated NA has a reduced immunogenicity in mice, as the variable and immunodominant surface loops on top of the NA molecule are masked by the N-glycans (Martinet et al. *Eur J. Biochem.* 247: 332-338, 1997).

Therefore, it is desirable to genetically engineer methy lotrophic yeast strains in which glycosylation of proteins can be manipulated and from which recombinant proteins can be produced that would not be compromised in structure or function by large N-glycan side chains.

SUMMARY OF THE INVENTION

The present invention is directed to methods and vectors useful for genetically modifying the glycosylation process in methy lotrophic yeast strains to produce glycoproteins with reduced glycosylation. Methy lotrophic yeast strains generated using the present methods and vectors, as well as glycoproteins produced from such genetically modified strains are also provided.

In one embodiment, the present invention provides vectors useful for making genetically engineered methy lotrophic yeast strains which are capable of producing glycoproteins with reduced glycosylation.

In one aspect, the present invention provides "knock-in" vectors which are capable of expressing in a methy lotrophic yeast strain one or more proteins whose enzymatic activities lead to a reduction of glycosylation in glycoproteins produced by the methy lotrophic yeast strain.

In a preferred embodiment, the knock-in vectors of the present invention include a nucleotide sequence coding for an α-1,2-mannosidase or a functional part thereof and are capable of expressing the α-1,2-mannosidase or the functional part in a methy lotrophic yeast strain. A preferred nucleotide sequence is a nucleotide sequence encoding the α-1,2-mannosidase of a fungal species, and more preferably, *Trichoderma reesei*. Preferably, the α-1,2-mannosidase expression vector is engineered such that the α-1,2-mannosidase or a functional part thereof expressed from the vector includes an ER-retention signal. A preferred ER-retention signal is HDEL (SEQ ID NO: 1). The α-1,2-mannosidase coding sequence can be operable linked to a constitutive or inducible promoter, and a 3' termination sequence. The vectors can be integrative vectors or replicative vectors. Particularly preferred α-1,2-mannosidase expression vectors include pGAPZMFManHDEL, pGAPZMFManMycHDEL,

pPICZBMFManMycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9 mMycManHDEL and pGAPZmMycManHDEL.

In another preferred embodiment, the knock-in vectors of the present invention include a sequence coding for a glucosidase II or a functional part thereof and are capable of expressing the glucosidase II or the functional part in a methylotrophic yeast strain. A preferred nucleotide sequence is a nucleotide sequence encoding the glucosidase II of a fungal species, and more preferably, *Saccharomyces cerevisiae*. Preferably, the glucosidase II expression vector is engineered such that the glucosidase II or a functional part thereof expressed from the vector includes an ER-retention signal. A preferred ER-retention signal is HDEL (SEQ ID NO: 1). The glucosidase II coding sequence can be operably linked to a constitutive or inducible promoter, and a 3' termination sequence. The vectors can be integrative vectors or replicative vectors. Particularly preferred glucosidase II expression vectors include pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, pGAPADEglsII, pPICADEglsII, pAOX2ADEglsII, pYPTIADeGlsII, pGAPZAgsIIHDEL and pGAPADEglsIIHDEL.

Expression vectors which include both of an α -1,2-mannosidase expression unit and a glucosidase II expression unit are also provided by the present invention.

In another aspect, the present invention provides "knock-out" vectors which, when introduced into a methylotrophic yeast strain, inactivate or disrupt a gene thereby facilitating the reduction in the glycosylation of glycoproteins produced in the methylotrophic yeast strain.

In one embodiment, the present invention provides a "knock-out" vector which, when introduced into a methylotrophic yeast strain, inactivates or disrupts the Och1 gene. A preferred Och1 knock-out vector is pBLURA5'PpOCH1.

Still another embodiment of the present invention provides vectors which include both a knock-in unit and a knock-out unit.

Furthermore, any of the knock-in or knock-out vectors of the present invention can also include a nucleotide sequence capable of expressing a heterologous protein of interest in a methylotrophic yeast.

Another embodiment of the present invention provides methods of modifying the glycosylation in a methylotrophic yeast by transforming the yeast with one or more vectors of the present invention.

Strains of a methylotrophic yeast which can be modified using the present methods include, but are not limited to, yeast strains capable of growth on methanol such as yeasts of the genera *Candida*, *Hansenula*, *Torulopsis*, and *Pichia*. Preferred methylotrophic yeasts are of the genus *Pichia*. Especially preferred are *Pichia pastoris* strains GS115 (NRRL Y-15851), GS190 (NRRL Y-18014), PPF1 (NRRL Y-18017), PPY120H, yGC4, and strains derived therefrom. Methylotrophic yeast strains which can be modified using the present methods also include those methylotrophic yeast strains which have been engineered to express one or more heterologous proteins of interest. The glycosylation on the heterologous proteins expressed from these previously genetically engineered strains can be reduced by transforming such strains with one or more of the vectors of the present invention.

Methylotrophic yeast strains which are modified by practicing the present methods are provided in another embodiment of the present invention.

A further aspect of the present invention is directed to methods of producing glycoproteins with a reduced glycosylation.

In accordance with such methods, a nucleotide sequence capable of expressing a glycoprotein can be introduced into a methylotrophic yeast strain which has previously been transformed with one or more of the vectors of the present invention. Alternatively, a methylotrophic yeast strain which has been genetically engineered to express a glycoprotein can be transformed with one or more of the vectors of the present invention. Moreover, if a methylotrophic yeast strain is not transformed with a nucleotide sequence encoding a glycoprotein of interest or any of the vectors of the present invention, such yeast strain can be transformed, either consecutively or simultaneously, with both a nucleotide sequence capable of expressing the glycoprotein and one or more vectors of the present invention. Additionally, a methylotrophic yeast strain can be transformed with one or more of the present knock-in and/or knock-out vectors which also include a nucleotide sequence capable of expressing a glycoprotein in the methylotrophic yeast strain.

Glycoproteins products produced by using the methods of the present invention, i.e., glycoproteins with reduced N-glycosylation, are also part of the present invention.

Kits which include one or more of the vectors of the present invention, or one or more strains modified to produce glycoproteins with reduced glycosylation, are also provided.

25

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts vectors carrying an HDEL (SEQ ID NO: 1)-tagged *Trichoderma reesei* α -1,2-mannosidase expression cassette and describes the way in which these vectors were constructed according to methods known in the art. Abbreviations used throughout construction schemes: 5' AOX1 or AOX1 P: *Pichia pastoris* AOX1 promoter sequence; Amp R: ampicillin resistance gene; ColE1: ColE1 origin of replication; 3'AOX1: 3' sequences of the *Pichia pastoris* AOX1 gene; HIS4: HIS4 gene of *Pichia pastoris*. AOX TT: transcription terminator sequence of the *Pichia pastoris* AOX1 gene; ORF: open reading frame; S: secretion signal; P TEF1: the promoter sequence of the *Saccharomyces cerevisiae* transcription elongation factor 1 gene; P EM7: synthetic constitutive prokaryotic promoter EM7; Zeocin: Zeocin resistance gene; CYC1 TT: 3' end of the *S. cerevisiae* CYC1 gene; GAP: promoter sequence of the *Pichia pastoris* glyceraldehyde-3-phosphate dehydrogenase gene; PpURA3: *Pichia pastoris* URA3 gene. As can be seen in this figure, the *Trichoderma reesei* α -1,2-mannosidase was operably linked to the coding sequence for the *S. cerevisiae* α -mating factor secretion signal sequence and further operably linked at the 3' terminus of the coding sequence to the coding sequence for an HDEL (SEQ ID NO: 1) peptide. The whole fusion construct was operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9MFManHDEL) or to the *P. pastoris* GAP promoter (in pGAPZMFManHDEL).

FIG. 2 depicts vectors carrying an HDEL (SEQ ID NO: 1)-tagged *Mus musculus* α -1,2-mannosidase IB expression cassette and describes the way in which these vectors were constructed according to methods known in the art. As can be seen in this figure, the catalytic domain of the *Mus musculus* α -1,2-mannosidase IB was operably linked to the coding sequence for the *S. cerevisiae* α -mating factor secretion signal sequence and further operably linked at the 3' terminus of the coding sequence to the coding sequence for an HDEL (SEQ ID NO: 1) peptide. The whole fusion construct was operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9 mManHDEL) or to the *P. pastoris* GAP promoter (in pGAPZmManHDEL). Furthermore, variants of the expression cassette were made in which the coding sequence for a

cMyc epitope tag was inserted between the coding sequence for the *S. cerevisiae* α-Mating Factor secretion signal sequence and the coding sequence for the catalytic domain of the *Mus musculus* α-1,2-mannosidase IB. This expression cassette was also operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9 mMycManHDEL) or to the *P. pastoris* GAP promotor (in pGAPZmMycManHDEL).

FIG. 3 depicts vectors carrying a MycHDEL tagged *Trichoderma reesei* α-1,2-mannosidase and the way in which these vectors were obtained. The resulting fusion construction was again operably linked to either the *P. pastoris* AOX1 promoter (in pPICZBMFManMycHDEL) or to the *P. pastoris* GAP promotor (in pGAPZMFManMycHDEL).

FIG. 4 demonstrates the intracellular localization of the MycHDEL-tagged *Trichoderma reesei* α-1,2-mannosidase and indicates ER-targeting by immunofluorescence analysis. Panel A Western blotting. Yeast strains were grown in 10 ml YPG cultures to an OD₆₀₀=10, diluted fivefold and grown in YPM for 48 h. 1/50th of the culture medium and 1/5th of the cells were analysed by SDS-PAGE and Western blotting with the mouse monoclonal 9E10 anti-Myc antibody. The position of molecular weight marker proteins are indicated with arrows. Lanes 1-5: cellular lysates. 1,2: pGAPZMFManMycHDEL transformants. 3: non-transformed PPY12OH (negative control). 4,5: pPICZBMFManMycHDEL transformants. Lanes 6-10: culture media. 6: non transformed PPY12OH (negative control). 7,8: pGAPZMFManMycHDEL transformants. 9,10: pPICZBMFManMycHDEL transformants. Panel B Immunofluorescence microscopy. 1: phase contrast image of a *P. pastoris* cell (strain PPY12OH transformed with pGAPZMFManHDEL) at 1000× magnification. The nucleus is visible as an ellipse in the lower right quadrant of the cell. 2: same cell as in 1, but in fluorescence microscopy mode to show localization of the *T. reesei* mannosidase-Myc-HDEL protein. The protein is mainly localized in a circular distribution around the nucleus (nuclear envelope), which is typical for an endoplasmic reticulum steady-state distribution. 3: phase contrast image of a *P. pastoris* cell (strain PPY12OH transformed with pGAPZMFManHDEL) at 1000× magnification. 4: same cell in fluorescence microscopy to show localization of the Golgi marker protein OCH1-HA in *P. pastoris* strain PPY12OH. The dot-like distribution throughout the cytoplasm, with 3-4 dots per cell is typical for cis-Golgi distribution in *P. pastoris*.

FIG. 5 depicts the co-sedimentation of mannosidase-MycHDEL with Protein Disulfide Isomerase in sucrose density gradient centrifugation. The top panel shows the distribution over the different fractions of the sucrose gradient of the OCH1-HA Golgi marker protein. The middle panel shows this distribution for the Protein Disulfide Isomerase endoplasmic reticulum marker protein. Finally, the bottom panel shows the distribution of the MycHDEL-tagged *Trichoderma reesei* α-1,2-mannosidase over the same fractions. It is concluded that the mannosidase-MycHDEL almost exactly matches the distribution of the ER marker PDI and thus mainly resides in the ER of the *Pichia pastoris* yeast cells.

FIG. 6 depicts the N-glycan analysis of *Trypanosoma cruzi* trans-sialidase coexpressed with *Trichoderma reesei* mannosidase-HDEL. Panel A: malto-oligosaccharide size reference ladder. Sizes of the glycans are expressed in Glucose Units (GU) by comparison of their electrophoretic mobility to the mobility of these malto-oligosaccharides. Panel B: N-glycans derived from recombinant *Trypanosoma cruzi* trans-sialidase expressed in *Pichia pastoris*. The peak at GU=9,2 corresponds to Man₈GlcNAc₂. Panel C: same analytes as panel 2, but after overnight treatment with 3 U/ml purified recombinant *T. reesei* α-1,2-mannosidase. Panel D: N-gly-

cans derived from recombinant trans-sialidase co-expressed in *Pichia pastoris* with *T. reesei* mannosidase-HDEL (under control of the GAP promotor). The peak at GU=7,6 corresponds to the Man₅GlcNAc₂ peak in the profile of RNase B (Panel F). Panel E: same analytes as panel D, but after overnight treatment with 3 mU/ml purified recombinant *T. reesei* α-1,2-mannosidase. Panel F: N-glycans derived from bovine RNase B. These glycans consist of Man₅GlcNAc₂ to Man₈GlcNAc₂. Different isomers are resolved, accounting for the number of peaks for Man₇GlcNAc₂.

FIG. 7 depicts the processing of influenza haemagglutinin N-glycans by HDEL (SEQ ID NO: 1)-tagged *Trichoderma reesei* α-1,2-mannosidase and the HDEL (SEQ ID NO: 1)-tagged catalytic domain of murine α-1,2-mannosidase IB. The Man₅GlcNAc₂ reference oligosaccharide runs at scan 1850 in this analysis (not shown). Panel 1: malto-oligosaccharide size reference ladder. Panel 2: N-glycans derived from recombinant influenza haemagglutinin expressed in *Pichia pastoris*. The peak at scan 2250 corresponds to Man₉GlcNAc₂. Panel 3: N-glycans derived from recombinant haemagglutinin co-expressed in *Pichia pastoris* with *T. reesei* mannosidase-HDEL (under control of the GAP promotor). The peak at scan 1950 corresponds to Man₆GlcNAc₂. Panel 4: Same analytes as for panel 3, but after overnight treatment with 30 mU purified recombinant *T. reesei* 1,2-mannosidase. Panel 5: N-glycans derived from recombinant haemagglutinin co-expressed in *Pichia pastoris* with mouse mannosidase IB-HDEL (under control of the GAP promotor). Panel 6: same analytes as for panel 5, but after overnight treatment with 30 mU purified recombinant *T. reesei* α-1,2-mannosidase.

FIG. 8 graphically depicts vector pBLURA5'PpOCH1 and the way in which it was constructed.

FIG. 9 depicts the scheme for disrupting the *Pichia pastoris* OCH1 gene by single homologous recombination using pBLURA5'PpOCH1.

FIG. 10 depicts the cell wall glycoprotein N-glycan analysis of the Och1-inactivated clone and three clones derived from this Och1-inactivated clone by transformation with pGAPZMFManHDEL. Panel 1 shows the analysis of a mixture of malto-oligosaccharides, the degree of polymerisation of which is given by the numbers on the very top of the figure. This analysis serves as a size reference for the other panels. On the vertical axis of all panels, peak intensity in relative fluorescence units is given. Panel 2-6: analysis of the cell wall glycoprotein N-glycans of the following strains: Panel 2, non-engineered *P. pastoris* strain yGC4; Panel 3, yGC4 transformed with pBLURA5'PpOch1; 4-6, three clones of the strain of Panel 3, supplementarily transformed with pGAPZMFManHDEL. Panel 7: the N-glycans derived from bovine RNaseB, consisting of a mixture of Man₅₋₉GlcNAc₂. As can be seen from comparison between panel 2 and 3 and reference to panel 7, transformation with pBLURA5'PpOch1 leads to a strongly increased abundance of the Man₈GlcNAc₂ substrate N-glycan (named peak 1 in Panel 2) of OCH1p. Peak 2 represents the Man₉GlcNAc₂ product of OCH1p. Furthermore, upon supplementary transformation of pGAPZMFManHDEL, the major glycan on the cell wall glycoproteins of three independent clones is the Man₅GlcNAc₂ end product (peak 3 in panel 4) of *T. reesei* α-1,2-mannosidase digestion of the Man₈GlcNAc₂ substrate.

FIG. 11 depicts the analysis of exactly the same glycan mixtures as in FIG. 10, but after an in vitro digest with 3 mU/ml purified *Trichoderma reesei* α-1,2-mannosidase, overnight in 20 mM sodium acetate pH=5.0. Axis assignment is the same as in FIG. 10. More Man₅GlcNAc₂ is formed in the pBLURA5'PpOch1 transformed strain (Panel 3) than in

the parent strain (Panel 2). Peaks in all panels before scan 3900 come from contaminants and should be ignored in the analysis.

FIG. 12 depicts the expression vector pGAPZAGLSII (SEQ ID NO: 18). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7: synthetic prokaryotic promoter. Zeocin: zeocine resistance marker gene. CYC1 TT: transcription terminator of *S. cerevisiae* cytochrome C1 gene. ColE1: bacterial origin of replication. GAP: promotor of the *P. pastoris* GAP gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

FIG. 13 depicts the expression vector pAOX2ZAGLSII (SEQ ID NO: 16). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7: synthetic prokaryotic promoter. Zeocin: zeocine resistance marker gene. CYC1 TT: transcription terminator of *S. cerevisiae* cytochrome C1 gene. Col E1: bacterial origin of replication. AOX2 P: promotor of the *P. pastoris* AOX2 gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

FIG. 14 depicts the expression vector pPICZAGLSII (SEQ ID NO: 20). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7: synthetic prokaryotic promoter. Zeocin: zeocine resistance marker gene. CYC1 TT: transcription terminator of *S. cerevisiae* cytochrome C1 gene. Col E1: origin of replication. AOX1 P: promotor of the *P. pastoris* AOX1 gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

FIG. 15 depicts the expression vector pYPT1ZAGLSII (SEQ ID NO: 22). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7: synthetic prokaryotic promoter. Zeocin: zeocine resistance marker gene. CYC1 TT: transcription terminator of *S. cerevisiae* cytochrome C1 gene. Col E1: origin of replication. P YPT1: promotor of the *P. pastoris* YPT1 gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene.

FIG. 16 depicts the expression vector pGAPADE1glsII (SEQ ID NO: 19). Amp R: Ampillicin resistance marker gene. ADE1: *P. pastoris* ADE1 selection marker gene. GAP: promotor of the *P. Pastoris* GAP gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

FIG. 17 depicts the expression vector pAOX2ADE1glsII (SEQ ID NO: 17). Amp R: Ampillicin resistance marker gene. ADE1: *P. pastoris* ADE1 selection marker gene. AOX2 P: promotor of the *P. pastoris* AOX2 gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene.

FIG. 18 depicts the expression vector pPICADE1glsII (SEQ ID NO: 21). Amp R: Ampillicin resistance marker gene. ADE1: *P. pastoris* ADE1 selection marker gene. AOX1 P: promotor of the *P. pastoris* AOX1 gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene.

FIG. 19 depicts the expression vector pYPT1ADE1glsII (SEQ ID NO: 23). Amp R: Ampillicin resistance marker gene. ADE1: *P. pastoris* ADE1 selection marker gene. P YPT1: promotor of the *P. pastoris* YPT1 gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene.

FIG. 20 depicts the expression vector pGAPZAglsIIHDEL (SEQ ID NO: 24). Amp R: Ampillicin resistance marker gene. ADE1: *P. pastoris* ADE1 selection marker gene. GAP: pro-

motor of the *P. pastoris* GAP gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene.

FIG. 21 depicts the expression vector pGAPADE1glsIIHDEL (SEQ ID NO: 25). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7: synthetic prokaryotic promoter. Zeocin: zeocine resistance marker gene. CYC1 TT: transcription terminator of *S. cerevisiae* cytochrome C1 gene. Col E1: bacterial origin of replication. GAP: promotor of the *P. pastoris* GAP gene. GLS2: *S. cerevisiae* glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene.

FIG. 22 depicts the test of the GLSII activity assay using a commercially available yeast alpha-glucosidase (Sigma: Cat. No. G-5003). The assay mixture contains phosphate-citrate buffer pH 6.8, mannose, 2-deoxy-D-glucose, the substrate 4-methylumbelliferyl-alpha-D-glucopyranoside and alpha-glucosidase from Sigma. 1: assay mixture illuminated with UV-light after overnight incubation at 37° C.; 2: same as 1, but this time, the assay mixture lacks the alpha-glucosidase; 3: same as 1, but this time, the assay mixture lacks the substrate.

FIG. 23 depicts the results of the activity of recombinantly expressed GLSII from *Pichia pastoris*. All assay mixtures were incubated overnight at 37° C. and afterwards illuminated with UV-light. 1: assay with yeast alpha-glucosidase (Sigma: Cat. No. G-5003); 2: assay with the purified medium of strain 18 (PPY12-OH transformed with pGAPZAGLSII); 3: assay with purified medium of the WT PPY12-OH strain; 4: assay with the purified medium of strain H3 (PPY12-OH transformed with pGAPZAglsIIHDEL).

DETAILED DESCRIPTION OF THE INVENTION

It has been established that the majority of N-glycans on glycoproteins leaving the endoplasmic reticulum (ER) of *Pichia* have the core $\text{Man}_8\text{GlcNAc}_2$ oligosaccharide structure. After the proteins are transported from the ER to the Golgi apparatus, additional mannose residues are added to this core sugar moiety by different mannosyltransferases, resulting in glycoproteins with large mannose side chains. Such hyperglycosylation of recombinant glycoproteins is undesirable in many instances. Accordingly, the present invention provides methods and vectors for genetically modifying methylotrophic yeast strains to produce glycoproteins with reduced glycosylation. Methylotrophic yeast strains generated using the present methods and vectors, as well as glycoproteins produced from such genetically modified strains are also provided.

In one embodiment, the present invention provides vectors useful for genetically modifying methylotrophic yeast strains to produce glycoproteins with reduced glycosylation.

In one aspect, the present invention provides “knock-in” vectors which are capable of expressing in a methylotrophic yeast strain one or more proteins whose enzymatic activities lead to a reduction of glycosylation in glycoproteins produced by the methylotrophic yeast strain. According to the present invention, such proteins include, e.g., an α -1,2-mannosidase, a glucosidase II, or functional parts thereof.

In a preferred embodiment, the vectors of the present invention include a sequence coding for an α -1,2-mannosidase or a functional part thereof and are capable of expressing the α -1,2-mannosidase or the functional part in a methylotrophic yeast strain.

An α -1,2-mannosidase cleaves the α -1,2-linked mannose residues at the non-reducing ends of $\text{Man}_8\text{GlcNAc}_2$, and converts this core oligosaccharide on glycoproteins to

$\text{Man}_5\text{GlcNAc}_2$. In vitro, $\text{Man}_5\text{GlcNAc}_2$ is a very poor substrate for any *Pichia* Golgi mannosyltransferase, i.e., mannose residues can not be added to this sugar structure. On the other hand, $\text{Man}_5\text{GlcNAc}_2$ is the acceptor substrate for the mammalian N-acetylglucosaminyl-transferase I and is an intermediate for the hybrid- and complex-type sugar chains characteristic of mammalian glycoproteins. Thus, by way of introducing an α -1,2-mannosidase into methylotrophic yeasts such as *Pichia*, glycoproteins with reduced mannose content can be produced.

According to the present invention, the nucleotide sequence encoding an α -1,2-mannosidase for use in the expression vector of the present invention can derive from any species. A number of α -1,2-mannosidase genes have been cloned and are available to those skilled in the art, including mammalian genes encoding, e.g., a murine α -1,2-mannosidase (Herscovics et al. *J. Biol. Chem.* 269: 9864-9871, 1994), a rabbit α -1,2-mannosidase (Lal et al. *J. Biol. Chem.* 269: 9872-9881, 1994) or a human α -1,2-mannosidase (Tremblay et al. *Glycobiology* 8: 585-595, 1998), as well as fungal genes encoding, e.g., an *Aspergillus* α -1,2-mannosidase (*msdS* gene), a *Trichoderma reesei* α -1,2-mannosidase (Maras et al. *J. Biotechnol.* 77: 255-263, 2000), or a *Saccharomyces cerevisiae* α -1,2-mannosidase. Protein sequence analysis has revealed a high degree of conservation among the eukaryotic α -1,2-mannosidases identified so far.

Preferably, the nucleotide sequence for use in the present vectors encodes a fungal α -1,2-mannosidase, more preferably, a *Trichoderma reesei* α -1,2-mannosidase, and more particularly, the *Trichoderma reesei* α -1,2-mannosidase described by Maras et al. *J. Biotechnol.* 77: 255-63 (2000).

According to the present invention, the nucleotide sequence can also code for only a functional part of an α -1,2-mannosidase.

By "functional part" is meant a polypeptide fragment of an α -1,2-mannosidase which substantially retains the enzymatic activity of the full-length protein. By "substantially" is meant at least about 40%, or preferably, at least 50% or more of the enzymatic activity of the full-length α -1,2-mannosidase is retained. For example, as illustrated by the present invention, the catalytic domain of the murine α -1,2-mannosidase IB constitutes a "functional part" of the murine α -1,2-mannosidase IB. Those skilled in the art can readily identify and make functional parts of an α -1,2-mannosidase using a combination of techniques known in the art. Predictions of the portions of an α -1,2-mannosidase essential to or sufficient to confer the enzymatic activity can be made based on analysis of the protein sequence. The activity of a portion of an α -1,2-mannosidase of interest, expressed and purified from an appropriate expression system, can be verified using in vitro or in vivo assays described hereinbelow.

In accordance with the present invention, an α -1,2-mannosidase or a functional part thereof expressed in a methylotrophic yeast strain preferably is targeted to a site in the secretory pathway where $\text{Man}_8\text{GlcNAc}_2$ (the substrate of α -1,2-mannosidase) is already formed on a glycoprotein, but has not reached a Golgi glycosyltransferase which elongates the sugar chain with additional mannose residues.

Accordingly, in a preferred embodiment of the present invention, the α -1,2-mannosidase expression vector is engineered as such that the α -1,2-mannosidase or a functional part thereof expressed from the vector includes an ER-retention signal.

"An ER retention signal" refers to a peptide sequence which directs a protein having such peptide sequence to be

transported to and retained in the ER. Such ER retention sequences are often found in proteins that reside and function in the ER.

Multiple choices of ER retention signals are available to those skilled in the art, e.g., the first 21 amino acid residues of the *S. cerevisiae* ER protein MNS1 (Martinet et al. *Biotechnology Letters* 20: 1171-1177, 1998). A preferred ER retention signal for use in the present invention is peptide HDEL (SEQ ID NO: 1). The HDEL (SEQ ID NO: 1) peptide sequence, found in the C-terminus of a number of yeast proteins, acts as a retention/retrieval signal for the ER (Pelham *EMBO J.* 7: 913-918, 1988). Proteins with an HDEL (SEQ ID NO: 1) sequence are bound by a membrane-bound receptor (Erd2p) and then enter a retrograde transport pathway for return to the ER from the Golgi apparatus.

According to the present invention, an ER retention signal can be placed anywhere in the protein sequence of an α -1,2-mannosidase, but preferably at the C-terminus of the α -1,2-mannosidase.

The α -1,2-mannosidase for use in the present invention can be further modified, e.g., by insertion of an epitope tag to which antibodies are available, such as Myc, HA, FLAG and His6 tags well-known in the art. An epitope-tagged α -1,2-mannosidase can be conveniently purified, or monitored for both expression and intracellular localization.

An ER retention signal and an epitope tag can be readily introduced into a protein of interest by inserting nucleotide sequences coding for such signal or tag into the nucleotide sequence encoding the protein of interest, using any of the molecular biology techniques known in the art.

In another preferred embodiment, the vectors of the present invention include a sequence coding for a glucosidase II or a functional part thereof and are capable of expressing the glucosidase II or the functional part in the methylotrophic yeast strain.

It has been established that the initial N-linked oligosaccharide ($\text{Glc}_3\text{Man}_9\text{GlcNAc}_2$), transferred in the ER onto a protein, is cleaved in the ER by specific glucosidases to remove the glucose residues, and by a mannosidase to remove one specific α -1,2-linked mannose. It has been observed by the present inventors that some recombinant proteins expressed in *Pichia* have residual glucose residues on the sugar moiety when such proteins leave the ER for the Golgi apparatus. The residual glucose molecules present on the sugar structure prevent the complete digestion of the sugar moiety by an α -1,2-mannosidase, and the introduction of an exogenous glucosidase can facilitate the removal of these glucose residues.

According to the present invention, the nucleotide sequence encoding a glucosidase II can derive from any species. Glucosidase II genes have been cloned from a number of mammalian species including rat, mouse, pig and human. The glucosidase II protein from these mammalian species consists of an alpha and a beta subunit. The alpha subunit is about 110 kDa and contains the catalytic activity of the enzyme, while the beta subunit has a C-terminal HDEL (SEQ ID NO: 1) ER-retention sequence and is believed to be important for the ER localization of the enzyme. The glucosidase II gene from *S. cerevisiae* has also been cloned (ORF YBR229c, located on chromosome II). This gene encodes a protein of about 110 kDa, which shows a high degree of homology to the mammalian alpha subunits.

A preferred glucosidase II gene for use in the present vectors is from a fungal species such as *Pichia pastoris* and *S. cerevisiae*. An example of a fungal glucosidase II gene is the *S. cerevisiae* glucosidase II alpha subunit gene.

11

According to the present invention, the nucleotide sequence can also encode only a functional part of a glucosidase II. By "functional part" is meant a polypeptide fragment of a glucosidase II which substantially retains the enzymatic activity of the full-length protein. By "substantially" is meant at least about 40%, or preferably, at least 50% or more of the enzymatic activity of the full-length glucosidase II is retained. Functional parts of a glucosidase II can be identified and made by those skilled in the art using a variety of techniques known in the art.

In a preferred embodiment of the present invention, the glucosidase II protein is engineered to include an ER retention signal such that the protein expressed in a methylotrophic yeast strain is targeted to the ER and retains therein for function. ER retention signals are as described hereinabove, e.g., the HDEL (SEQ ID NO: 1) peptide sequence.

The glucosidase II for use in the present invention can be further modified, e.g., by insertion of an epitope tag to which antibodies are available, such as Myc, HA, FLAG, and His6 tag, which are well-known in the art.

According to the present invention, the "knock-in" vectors can include either or both of an α -1,2-mannosidase coding sequence and a glucosidase II coding sequence.

Further according to the present invention, the nucleotide sequence coding for the enzyme to be expressed (e.g., an α -1,2-mannosidase or a functional part thereof, or a glucosidase II or a functional part thereof) can be placed in an operable linkage to a promoter and a 3' termination sequence.

Promoters appropriate for expression of a protein in a methylotrophic yeast can include both constitutive promoters and inducible promoters. Constitutive promoters include e.g., the *Pichia pastoris* glyceraldehyde-3-phosphate dehydrogenase promoter ("the GAP promoter"). Examples of inducible promoters include, e.g., the *Pichia pastoris* alcohol oxidase I promoter ("the AOXI promoter") (U.S. Pat. No. 4,855,231), or the *Pichia pastoris* formaldehyde dehydrogenase promoter ("the FLD promoter") (Shen et al. Gene 216: 93-102, 1998).

3' termination sequences are sequences 3' to the stop codon of a structural gene which function to stabilize the mRNA transcription product of the gene to which the sequence is operably linked, such as sequences which elicit polyadenylation. 3' termination sequences can be obtained from *Pichia* or other methylotrophic yeast. Examples of *Pichia pastoris* 3' termination sequences useful for the practice of the present invention include termination sequences from the AOX1 gene, p40 gene, HIS4 gene and FLD1 gene.

The vectors of the present invention preferably contain a selectable marker gene. The selectable marker may be any gene which confers a selectable phenotype upon a methylotrophic yeast strain and allows transformed cells to be identified and selected from untransformed cells. The selectable marker system may include an auxotrophic mutant methylotrophic yeast strain and a wild type gene which complements the host's defect. Examples of such systems include the *Saccharomyces cerevisiae* or *Pichia pastoris* HIS4 gene which may be used to complement his4 *Pichia* strains, or the *S. cerevisiae* or *Pichia pastoris* ARG4 gene which may be used to complement *Pichia pastoris* arg mutants. Other selectable marker genes which function in *Pichia pastoris* include the Zeo^R gene, the G418^R gene, and the like.

The vectors of the present invention can also include an autonomous replication sequence (ARS). For example, U.S. Pat. No. 4,837,148 describes autonomous replication sequences which provide a suitable means for maintaining plasmids in *Pichia pastoris*. The disclosure of U.S. Pat. No. 4,837,148 is incorporated herein by reference.

12

The vectors can also contain selectable marker genes which function in bacteria, as well as sequences responsible for replication and extrachromosomal maintenance in bacteria. Examples of bacterial selectable marker genes include 5 ampicillin resistance (Amp^R), tetracycline resistance (Tet^R), neomycin resistance, hygromycin resistance, and zeocin resistance (Zeo^R) genes.

According to the present invention, the nucleotide sequence encoding the protein to be expressed in a methylotrophic yeast can be placed in an integrative vector or a replicative vector (such as a replicating circular plasmid).

Integrative vectors are disclosed, e.g., in U.S. Pat. No. 4,882,279 which is incorporated herein by reference. Integrative vectors generally include a serially arranged sequence of at least a first insertable DNA fragment, a selectable marker gene, and a second insertable DNA fragment. The first and second insertable DNA fragments are each about 200 nucleotides in length and have nucleotide sequences which are homologous to portions of the genomic DNA of the species to be transformed. A nucleotide sequence containing a structural gene of interest for expression is inserted in this vector between the first and second insertable DNA fragments whether before or after the marker gene. Integrative vectors can be linearized prior to yeast transformation to facilitate the integration of the nucleotide sequence of interest into the host cell genome.

Replicative and integrative vectors carrying either or both of an α -1,2-mannosidase coding sequence or a glucosidase II coding sequence can be constructed by standard techniques known to one of ordinary skill in the art and found, for example, in Sambrook et al. (1989) in Molecular Cloning: A Laboratory Manual, or any of a myriad of laboratory manuals on recombinant DNA technology that are widely available.

Preferred vectors of the present invention carrying an α -1,2-mannosidase expression sequence include pGAPZMF-ManHDEL, pGAPZMFMManMycHDEL, pPICZBMFMan-MycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9 mMycManHDEL and 40 pGAPZmMycManHDEL, which are further described in the Examples hereinbelow.

Preferred vectors of the present invention carrying a glucosidase II expression sequence include pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, 45 pGAPADE1glsII, pPICADE1glsII, pAOX2ADE1glsII, pYPTIADE1glsII, pGAP

ZAGlsIIHDEL and pGAPADE1glsIIHDEL, which are further described in the Examples hereinbelow.

In another aspect, the present invention provides "knock-out" vectors which, when introduced into a methylotrophic yeast strain, inactivate or disrupt a gene thereby facilitating the reduction in the glycosylation of glycoproteins produced in the methylotrophic yeast strain.

In one embodiment, the present invention provides a 55 "knock-out" vector which, when introduced into a methylotrophic yeast strain, inactivates or disrupts the Och1 gene.

The *S. cerevisiae* OCH1 gene has been cloned (Nakayama et al. EMBO J. 11: 2511-2519, 1992). It encodes a membrane bound α -1,6-mannosyltransferase, localized in the early Golgi complex, that is functional in the initiation of α -1,6-polymannose outer chain addition to the N-linked core oligosaccharide (Man₅GlcNAc₂ and Man₈GlcNAc₂) (Nakanishi-Shindo et al. J. Biol. Chem. 268: 26338-26345, 1993).

A *Pichia* sequence has been described in Japanese Patent Application No. 07145005 that encodes a protein highly 65 homologous to the *S. cerevisiae* OCH1. For purpose of the present invention, this sequence is denoted herein as "the

Pichia OCH1 gene". Those skilled in the art can isolate the OCH1 genes from other methylotrophic yeasts using techniques well known in the art.

According to the present invention, a disruption in the OCH1 gene of a methylotrophic yeast can result in either the production of an inactive protein product or no product. The disruption may take the form of an insertion of a heterologous DNA sequence into the coding sequence and/or the deletion of some or all of the coding sequence. Gene disruptions can be generated by homologous recombination essentially as described by Rothstein (in *Methods in Enzymology*, Wu et al., eds., vol 101:202-211, 1983).

To disrupt the Och1 gene by homologous recombination, an Och1 knock-out vector can be constructed in such a way to include a selectable marker gene. The selectable marker gene is operably linked, at both 5' and 3' end, to portions of the Och1 gene of sufficient length to mediate homologous recombination. The selectable marker can be one of any number of genes which either complement host cell auxotrophy or provide antibiotic resistance, including URA3, LEU2 and HIS3 genes. Other suitable selectable markers include the CAT gene, which confers chloramphenicol resistance on yeast cells, or the lacZ gene, which results in blue colonies due to the expression of active β-galactosidase. Linearized DNA fragments of an Och1 knock-out vector are then introduced into host methylotrophic yeast cells using methods well known in the art. Integration of the linear fragments into the genome and the disruption of the Och1 gene can be determined based on the selection marker and can be verified by, for example, Southern Blot analysis.

Alternatively, an Och1 knock-out vector can be constructed in such a way to include a portion of the Och1 gene to be disrupted, which portion is devoid of any Och1 promoter sequence and encodes none or an inactive fragment of the Och1 protein. By "an inactive fragment", it is meant a fragment of the Och1 protein which has, preferably, less than about 10% and most preferably, about 0% of the activity of the full-length OCH1 protein. Such portion of the OCH1 gene is inserted in a vector in such a way that no known promoter sequence is operably linked to the OCH1 sequence, but that a stop codon and a transcription termination sequence are operably linked to the portion of the Och1 gene. This vector can be subsequently linearized in the portion of the OCH1 sequence and transformed into a methylotrophic yeast strain using any of the methods known in the art. By way of single homologous recombination, this linearized vector is then integrated in the OCH1 gene. Two Och1 sequences are produced in the chromosome as a result of the single homologous recombination. The first Och1 sequence is the portion of the Och1 gene from the vector, which is now under control of the OCH1 promoter of the host methylotrophic yeast, yet cannot produce an active OCH1 protein as such Och1 sequence codes for no or an inactive fragment of the OCH1 protein, as described hereinabove. The second Och1 sequence is a full OCH1 coding sequence, but is not operably linked to any known promoter sequence and thus, no active messenger is expected to be formed for synthesis of an active OCH1 protein. Preferably, an inactivating mutation is introduced in the OCH1 sequence, to the 5' end of the site of linearization of the vector and to the 3' end of the translation initiation codon of OCH1. By "inactivating mutation" it is meant a mutation introducing a stop codon, a frameshift mutation or any other mutation causing a disruption of the reading frame. Such mutation can be introduced into an Och1 sequence using any of the site directed mutagenesis methods known in the art. Such inactivating mutation ensures that no functional OCH1

protein can be formed even if there exist some promoter sequences 5' to the Och1 sequence in the knock-out vector.

A preferred Och1 knock-out vector of the present invention is pBLURA5'PpOCH1.

If desired, either or both of a mannosidase expression sequence and a glucosidase expression sequence can be carried on the same plasmid used to disrupt the OCH1 gene to create a "knock-in-and-knock-out" vector.

Additionally, any of the above-described vectors can further include a nucleotide sequence capable of expressing a glycoprotein of interest in a methylotrophic yeast strain.

Another aspect of the present invention is directed to methods of modifying methylotrophic yeast strains to reduce glycosylation on proteins produced by the methylotrophic yeast strains. In accordance with the present methods, methylotrophic yeast strains are modified by transforming into these yeast strains with one or more, i.e., at least one, knock-in and/or knock-out vectors of the present invention as described herein above.

Methylotrophic yeast strains which can be modified using the present methods include but are not limited to yeast capable of growth on methanol such as yeasts of the genera *Candida*, *Hansenula*, *Torulopsis*, and *Pichia*. A list of species which are exemplary of this class of yeasts can be found in C. Anthony (1982), *The Biochemistry of Methylotrophs*, 269. *Pichia pastoris*, *Pichia methanolica*, *Pichia anomola*, *Hansenula polymorpha* and *Candida boidinii* are examples of methylotrophic yeasts useful in the practice of the present invention. Preferred methylotrophic yeasts are of the genus *Pichia*. Especially preferred are *Pichia pastoris* strains GS115 (NRRL Y-15851); GS 190 (NRRL Y-18014) disclosed in U.S. Pat. No. 4,818,700; PPF1 (NRRL Y-18017) disclosed in U.S. Pat. No. 4,812,405; PPY120H and yGC4; as well as strains derived therefrom.

Methylotrophic yeast strains which can be modified using the present methods also include those methylotrophic yeast strains which have been genetically engineered to express one or more heterologous glycoproteins of interest. The glycosylation on the heterologous glycoproteins expressed from these previously engineered strains can be reduced by transforming such strains with one or more of the vectors of the present invention.

The vectors of the present invention can be introduced into the cells of a methylotrophic yeast strain using known methods such as the spheroplast technique, described by Cregg et al. 1985, or the whole-cell lithium chloride yeast transformation system, Ito et al. *Agric. Biol. Chem.* 48:341, modified for use in *Pichia* as described in EP 312,934. Other published methods useful for transformation of the plasmids or linear vectors include U.S. Pat. No. 4,929,555; Hinnen et al. *Proc. Nat. Acad. Sci. USA* 75:1929 (1978); Ito et al. *J. Bacteriol.* 153:163 (1983); U.S. Pat. No. 4,879,231; Sreekrishna et al. *Gene* 59:115 (1987). Electroporation and PEG1000 whole cell transformation procedures may also be used. Cregg and Russel *Methods in Molecular Biology: Pichia Protocols*, Chapter 3, Humana Press, Totowa, N.J., pp. 27-39 (1998).

Transformed yeast cells can be selected by using appropriate techniques including but not limited to culturing auxotrophic cells after transformation in the absence of the biochemical product required (due to the cell's auxotrophy), selection for and detection of a new phenotype, or culturing in the presence of an antibiotic which is toxic to the yeast in the absence of a resistance gene contained in the transformants. Transformants can also be selected and/or verified by integration of the expression cassette into the genome, which can be assessed by e.g., Southern Blot or PCR analysis.

15

In one embodiment, a methylotrophic yeast strain is transformed with a vector which includes a nucleotide sequence coding for an α -1,2-mannosidase or a functional part thereof. The nucleotide sequence is capable of expressing the α -1,2-mannosidase or the functional part in the methylotrophic yeast strain, and is, preferably, integrated into the genome of the methylotrophic yeast strain.

The expression of an α -1,2-mannosidase introduced in a methylotrophic yeast strain can be verified both at the mRNA level, e.g., by Northern Blot analysis, and at the protein level, e.g., by Western Blot analysis. The intracellular localization of the protein can be analyzed by using a variety of techniques, including subcellular fractionation and immunofluorescence experiments. An ER localization of an α -1,2-mannosidase can be determined by co-sedimentation of this enzyme with a known ER resident protein (e.g., Protein Disulfide Isomerase) in a subcellular fractionation experiment. An ER localization can also be determined by an immunofluorescence staining pattern characteristic of ER resident proteins, typically a perinuclear staining pattern.

To confirm that an α -1,2-mannosidase or a functional part thereof expressed in a methylotrophic yeast strain has the expected mannose-trimming activity, both *in vitro* and *in vivo* assays can be employed. Typically, an *in vitro* assay involves digestion of an *in vitro* synthesized substrate, e.g., $\text{Man}_8\text{GlcNAc}_2$, with the enzyme expressed and purified from a methylotrophic yeast strain, and assessing the ability of such enzyme to trim $\text{Man}_8\text{GlcNAc}_2$ to, e.g., $\text{Man}_5\text{GlcNAc}_2$. In *in vivo* assays, the α -1,2-mannosidase or a part thereof is co-expressed in a methylotrophic yeast with a glycoprotein known to be glycosylated with N-glycans bearing terminal α -1,2-linked mannose residues in such yeast. The enzymatic activity of such an α -1,2-mannosidase or a part thereof can be measured based on the reduction of the number of α -1,2-linked mannose residues in the structures of the N-glycans of the glycoprotein. In both *in vitro* and *in vivo* assays, the composition of a carbohydrate group can be determined using techniques that are well known in the art and are illustrated in the Examples hereinbelow.

In another embodiment, a methylotrophic yeast strain is transformed with a vector which includes a nucleotide sequence coding for a glucosidase II or a functional part thereof. The nucleotide sequence is capable of expressing the glucosidase II or the functional part in the methylotrophic yeast strain, and is, preferably, integrated into the genome of the methylotrophic yeast strain.

The enzymatic activity of a glucosidase II or a functional part thereof expressed in a transformed methylotrophic yeast strain can be assessed using a variety of assays. For example, methylotrophic yeast cells transformed with a sequence encoding a glucosidase II or a part thereof can be set to grow on solid medium containing a substrate of the glucosidase, e.g., 5-bromo-4-chloro-3-indolyl- α -D-glucopyranoside or 4-MU- α -D-Glc. When the enzyme is expressed by the *Pichia* and secreted extracellularly, the substrate is acted upon by the enzyme, giving rise to detectable signals around the colonies such as blue color or fluorescent glow. Alternatively, liquid culture medium containing the expressed protein molecules can be collected and incubated in test tubes with a substrate, e.g., p-nitrophenyl- α -D-glucopyranoside. The enzymatic activity can be determined by measuring the specific product released. Moreover, *in vivo* assays can be employed, where a glucosidase II is co-expressed in yeast with a glycoprotein known to be N-glycosylated with glucose residues, e.g., influenza neuraminidase. The enzymatic activity of the glucosidase II can be measured based on the reduction of the glucose content in the sugar chain(s) of the glycoprotein.

16

In still another embodiment of the present invention, a methylotrophic yeast strain is transformed with an Och1 knock-out vector. As a result of the transformation and integration of the vector, the genomic Och1 gene in the yeast strains is disrupted.

In a further embodiment of the present invention, a methylotrophic yeast strain is transformed with any combination of an α -1,2-mannosidase expression vector, a glucosidase II expression vector, and an Och1 knock-out vector. Such modification can be achieved by serial, consecutive transformations, i.e., introducing one vector at a time, or alternatively by co-transformation, i.e., introducing the vectors simultaneously.

The modified methylotrophic yeast strains described herein above can be further modified if desired. For example, additional disruption of genes encoding any other *Pichia* mannosyltransferases can be made. Genes encoding mammalian enzymes can also be introduced to produce glycoproteins having hybrid- or complex-type N-glycans, if desired.

Methylotrophic yeast strains which are modified by using the present methods, i.e., by transforming with one or more of the vectors of the present invention, form another embodiment of the present invention.

It should be understood that certain aspects of the present invention, especially the introduction of an intracellularly expressed α -1,2-mannosidase activity, are also useful to obtain a reduced glycosylation of the O-linked glycans on glycoproteins produced in a methylotrophic yeast, as it is known in the art that these O-linked glycans consist mainly of α -1,2-linked mannose residues. O-linked glycans as used herein refers to carbohydrate structures linked to serine or threonine residues of glycoproteins.

A further aspect of the invention is directed to methods of producing a glycoprotein with reduced glycosylation in a methylotrophic yeast, especially a glycoprotein heterologous to the methylotrophic yeast.

“A glycoprotein” as used herein refers to a protein which, in methylotrophic yeasts, is either glycosylated on one or more asparagines residues or on one or more serine or threonine residues, or on both asparagines and serine or threonine residues.

The term “reduced glycosylation” refers to a reduced size of the carbohydrate moiety on the glycoprotein, particularly with fewer mannose residues, when the glycoprotein is expressed in a methylotrophic yeast strain which has been modified in accordance with the present invention, as compared to a wild type, unmodified strain of the methylotrophic yeast.

In accordance with the present invention, the production of a glycoprotein of interest with reduced glycosylation can be achieved in a number of ways. A nucleotide sequence capable of expressing a glycoprotein can be introduced into a methylotrophic yeast strain which has been previously modified in accordance with the present invention, i.e., a strain transformed with one or more of the vectors of the present invention and capable of producing glycoproteins with reduced glycosylation. Alternatively, a methylotrophic yeast strain which has already been genetically engineered to express a glycoprotein can be transformed with one or more of the vectors of the present invention. Otherwise, if a methylotrophic yeast strain does not express a glycoprotein of interest, nor is the strain transformed with any of the vectors of the present invention, such yeast strain can be transformed, either consecutively or simultaneously, with both a nucleotide sequence capable of expressing the glycoprotein and one or more vectors of the present invention. Additionally, a methylotrophic yeast strain can be transformed with one or more of

17

the present knock-in and/or knock-out vectors which also include a nucleotide sequence capable of expressing a glycoprotein in the methylotrophic yeast strain.

The nucleotide sequence capable of expressing a glycoprotein in a methylotrophic yeast can be made to include from 5' to 3', a promoter, a sequence encoding the glycoprotein, and a 3' termination sequence. Promoters and 3' termination sequences which are suitable for expression of a glycoprotein can include any of those promoters and 3' termination sequences described hereinabove.

The nucleotide sequence for expression of a glycoprotein can include additional sequences, e.g., signal sequences coding for transit peptides when secretion of a protein product is desired. Such sequences are widely known, readily available and include *Saccharomyces cerevisiae* alpha mating factor prepro (*αmf*), *Pichia pastoris* acid phosphatase (PHO1) signal sequence and the like.

The nucleotide sequence for expression of a glycoprotein can be placed on a replicative vector or an integrative vector. The choice and construction of such vectors are as described hereinabove.

The nucleotide sequence capable of expressing a glycoprotein can be carried on the same replicative plasmid as a plasmid-borne α-1,2-mannosidase or glucosidase II expression unit. Alternatively, the nucleotide sequence containing the glycoprotein coding sequence is carried on a separate plasmid or integrated into the host genome.

Glycoproteins produced can be purified by conventional methods. Purification protocols can be determined by the nature of the specific protein to be purified. Such determination is within the ordinary level of skill in the art. For example, the cell culture medium is separated from the cells and the protein secreted from the cells can be isolated from the medium by routine isolation techniques such as precipitation, immunoabsorption, fractionation or a variety of chromatographic methods.

Glycoproteins which can be produced by the methods of the present invention include, e.g., *Bacillus amyloliquefaciens* α-amylase, *S. cerevisiae* invertase, *Trypanosoma cruzi* trans-sialidase, HIV envelope protein, influenza virus A haemagglutinin, influenza neuraminidase, Bovine herpes virus type-1 glycoprotein D, human angiostatin, human B7-1, B7-2 and B-7 receptor CTLA-4, human tissue factor, growth factors (e.g., platelet-derived growth factor), tissue plasminogen activator, plasminogen activator inhibitor-I, urokinase, human lysosomal proteins such as α-galactosidase, plasminogen, thrombin, factor XIII and immunoglobulins. For additional useful glycoproteins which can be expressed in the genetically engineered *Pichia* strains of the present invention, see Bretthauer and Castellino, *Biotechnol. Appl. Biochem.* 30: 193-200 (1999), and Kukuruzinska et al. *Ann Rev. Biochem.* 56: 915-44 (1987).

Glycoproteins produced by using the methods of the present invention, i.e., glycoproteins with reduced glycosylation, are also part of the present invention.

Still another aspect of the present invention provides kits which contain one or more of the knock-in vectors, knock-out vectors, or knock-in-and-knock-out vectors of the present invention described above. More particularly, a kit of the present invention contains a vector capable of expressing an α-mannosidase I in a methylotrophic yeast, a vector capable of expressing a glucosidase II in a methylotrophic yeast, a vector capable of disrupting the Och1 gene in a methylotrophic yeast, a vector capable of expressing both a glucosidase II and an α-mannosidase, a vector capable of

18

disrupting the Och1 gene and capable of expressing either or both of a glucosidase II and an α-mannosidase, or any combinations thereof.

The kit can also include a nucleotide sequence which encodes and is capable of expressing a heterologous glycoprotein of interest. Such nucleotide sequence can be provided in a separate vector or in the same vector which contains sequences for knocking-in or knocking out as described hereinabove.

In addition, the kit can include a plasmid vector in which a nucleotide sequence encoding a heterologous protein of interest can be subsequently inserted for transformation into and expression in a methylotrophic yeast. Alternatively, the knock-in or knock-out vectors in the kits have convenient cloning sites for insertion of a nucleotide sequence encoding a heterologous protein of interest.

The kit can also include a methylotrophic yeast strain which can be subsequently transformed with any of the knock-in, knock-out or knock-in-and-knock-out vectors described hereinabove. The kit can also include a methylotrophic yeast strain which has been transformed with one or more of the knock-in or knock-out vectors. Furthermore, the kit can include a methylotrophic yeast strain which has been transformed with a nucleotide sequence encoding and capable of expressing a heterologous glycoprotein of interest.

The present invention is further illustrated by the following examples.

Example 1

Introduction of α-1,2-Mannosidase to the ER-Golgi Border

1.1 Plasmids

	Plasmid	Promoter	Enzyme	Tag
30	pGAPZMFManHDEL	GAP	<i>T. reesei</i> α-1,2-mannosidase	—
40	pGAPZMFManMycHDEL	GAP	<i>T. reesei</i> α-1,2-mannosidase	Myc
45	pPICZBMFManMycHDEL	AOX1	<i>T. reesei</i> α-1,2-mannosidase	Myc
35	pGAPZMFmManHDEL	GAP	mouse mannosidase IB catalytic domain	—
50	pGAPZMFmMycManHDEL	GAP	mouse mannosidase IB catalytic domain	Myc

The *Trichoderma reesei* α-1,2-mannosidase gene has been isolated and described by Maras et al. (*J. Biotechnol.* 77: 255-263, 2000). The sequence of this gene is available at NCBI Genbank under Accession No. AF212153. A construction fragment was generated by PCR using the pPIC9MFmanase plasmid (same as pPP1MFmds1 described by Maras et al. (2000)) as the template and using the following oligonucleotide primers: 5'-GACTGGTTCCAATTGACAAGC-3' (SEQ ID NO: 2) and 5'-AGTCTAGATTACAACTCGTCGTGAGCAAGGTGGCCGCCCG TCG-3' (SEQ ID NO: 3). The resulting product contained the 3' end of the *Pichia pastoris* AOX1 promoter, the prepro-signal sequence of the *S. cerevisiae* α-mating factor, the open reading frame of the *Trichoderma reesei* α-1,2-mannosidase cloned in frame with the signal sequence, the coding sequence for HDEL (SEQ ID NO: 1), a stop codon and an Xba I restriction site. This fragment was digested with Eco RI and Xba I, removing the 5' sequences up to the mannosidase ORF, and then cloned into the vector pGAPZαA (Invitrogen, Baarn, The Netherlands) which had been digested with Eco

RI and Xba I, thus restoring the fusion with the *S. cerevisiae* α -mating factor signal sequence. The resulting plasmid was named pGAPZMFManHDEL and is graphically depicted in FIG. 1. The ORF sequence of the MFManHDEL fusion in pGAPZMFManHDEL is set forth in SEQ ID NO: 14.

In order to introduce the coding sequence for a c-Myc tag between the catalytic domain and the HDEL-signal (SEQ ID NO: 1), the 3' end of the ORF of *T. reesei* α -1,2-mannosidase was PCR-amplified using a sense primer 5'-CCATTGAG-GACGCATGCCGCC-3' (SEQ ID NO: 4) (containing an Sph I restriction site) and an antisense primer GTATCTA-GATTACAACACTCGTCGTGCAGATCCTCT-TCTGAGATGAGTTTGTTC AGCAAGGTGGCCGC-CCCCTCGTGATGAA (SEQ ID NO: 5) (containing the coding sequences of the c-Myc tag and the HDEL (SEQ ID NO: 1) signal, followed by a stop codon and an Xba I restriction site). The resulting PCR product was digested with Sph I and Xba I, purified by agarose gel electrophoresis and inserted into pGAPZMFManHDEL which had been cut with the same restriction enzymes, resulting in plasmid pGAPZMFManMycHDEL. To put the ORF of pGAPZMFManMycHDEL under the control of the inducible AOXI promoter, the entire ORF was liberated from pGAPZMFManMycHDEL with Bst BI and Xba I, and cloned in pPICZB (Invitrogen, Baarn, The Netherlands), resulting in pPICZBMFManMycHDEL.

Cloning of the mouse mannosidase IB catalytic domain with concomitant addition of the coding sequence for a C-terminal HDEL-tag (SEQ ID NO: 1) was done by PCR on a mouse cDNA library (mRNA isolated from the L929 cell line induced with cycloheximide and mouse Tumor Necrosis Factor. Average insert length of the cDNA library was 2000 bp). The PCR oligonucleotide primers used were: 5'AACTC-GAGATGGACTCTTCAAAACACAAACGC3' (SEQ ID NO: 6) and 5'TTGCAGCGCTTACAACCTCGTCGT-GTCGGACAGCAGGATTACCTGA3' (SEQ ID NO: 7). The product contained a 5' Xho I site and the coding sequence for C-terminal HDEL-site, followed by a stop codon and a Not I site at the 3' end. The product was cloned in pGAPZaA via the Xho I/Not I sites in the PCR product and the vector, resulting in an in frame fusion of the mouse mannosidase catalytic domain with the *S. cerevisiae* α -mating factor signal sequence. The sequence of the entire open reading frame generated is set forth in SEQ ID NO: 15.

1.2 Yeast Transformation and Genomic Integration

TABLE 2

Parental strain	DNA transformed
GS115 (his4)	pGAPZMFManHDEL pPIC9MFManHDEL pPIC9mManHDEL pPIC9mMycManHDEL pGAPZmManHDEL pGAPZmMycManHDEL pGAPZMFManHDEL pGAPZmManHDEL pGAPZmMycManHDEL
GS115 (his4 complemented by pPIC9InfluenzaHA)	
PPY120H (his4 complemented by pPIC9sOCH1)	pGAPZMFManMycHDEL pPICZBMFManMycHDEL
yGC4 (his4 arg1 ade2 ura3 complemented by pBLURAS5'PpOCH1)	pPIC9InfluenzaNeuraminidase pGAPZMFManHDEL pPIC9Glucoseoxidase

All transformations to *Pichia pastoris* were performed with electroporation according to the directions of Invitrogen. Transformants of vectors carrying the Zeocin resistance gene were selected on YPD containing 100 μ g/ml Zeocine (Invitrogen, Baarn, the Netherlands) and 1M sorbitol. Selection of

transformants of pPIC9 derivatives was done on minimal medium lacking histidine and containing 1M sorbitol. Genomic integration of the expression cassettes was verified using PCR on genomic DNA purified from the *Pichia* strains using the Yeast Miniprep method (Nucleon). In all cases concerning the *Trichoderma reesei* gene fusions, the primers used were the sense primer 5'-CCATTGAGGACGCATGC-CGCGCC-3' (SEQ ID NO: 8), which annealed to the 3' half of the mannosidase ORF, and the antisense primer 3' AOXI 5'-GCAAATGGCATTCTGACATCCT-3' (SEQ ID NO: 9), which annealed to the AOXI transcription terminator that was present in all our expression constructs. For the control of genomic integration of the mouse mannosidase transgenes, PCR was done using the sense primer 5'GAP 5'GTC-CCTATTCAATCAATTGAA3' (SEQ ID NO: 10, annealing to the GAP promoter or 5'AOXI 5'GACTGGTTCCAAT-TGACAAGC3' (SEQ ID NO: 11), annealing to AOXI promoter), and the antisense primer 3'AOXI (above). For the expression constructs containing a Myc tagged *Trichoderma reesei* α -1,2-mannosidase expression unit, further evidence for genomic integration was obtained using Southern Blotting with the entire MFManMycHDEL ORF (32 P labelled using HighPrime, Boehringer Mannheim) as a probe.

1.3 Expression of α -1,2-mannosidase

Expression of an α -1,2-Mannosidase in GS115 strains expressing influenza virus haemagglutinin was verified by qualitative Northern blot. Expression of an α -1,2-Mannosidase in PPY120H strains was verified by anti-Myc Western blot.

Qualitative Northern Blot—

Total RNA was purified from *Pichia* strains and the yield was determined spectrophotometrically. Northern blotting was performed according to standard procedures and an estimate of the quantity of RNA loaded was made using methylene blue staining of the blot, visualizing the rRNA bands. The blot was probed with a ClaI/NarI fragment of the mannosidase, labelled with 32 P using HighPrime (Boehringer Mannheim).

SDS-PAGE and Western Blotting—

Total yeast cell lysates were prepared by washing the cells twice with PBS, followed by boiling in 1 volume of 2x concentrated Laemmli loading buffer for 5 min. The lysate was spun briefly in a microcentrifuge prior to gel loading and only the supernatant was loaded. For the analysis of proteins secreted into the growth media, the proteins were precipitated from 200 μ l of these media using desoxycholate/trichloroacetic acid according to standard procedures. The pellet was redissolved in 2x concentrated Laemmli loading buffer and the solutions were pH-corrected using Tris. SDS-PAGE was performed and followed by semidry electroblotting to nitrocellulose membranes. For Western Blotting, the 9E10 anti-Myc and the anti-HA mouse monoclonals (Boehringer Mannheim) were used at a concentration of 1 μ g/ml, and the rabbit anti-PDI antiserum (Stressgen) was used at a dilution of 1/500. The secondary antibodies were goat anti-mouse IgG conjugated to alkaline phosphatase for the monoclonals and goat anti-rabbit IgG conjugated to peroxidase for the polyclonal (secondary antibodies from Sigma). Detection was performed using the NBT/BCIP system for alkaline phosphatase and the Renaissance substrate (NENBiosciences) for peroxidase. Imaging of the latter blot result was done on a Lumilager imaging device (Boehringer Mannheim).

The results shown in FIG. 4 indicated that the great majority of the HDEL (SEQ ID NO: 1)-tagged protein was retained intracellularly, both when expressed from the strong constitutive GAP promoter and when expressed from the strong inducible AOXI promoter.

21

1.4 Localization of α -1,2-Mannosidase

Isopycnic Sucrose Density Gradient Centrifugation—

To determine the localization of the HDEL (SEQ ID NO: 1)-tagged mannosidase, subcellular fractionation was carried out using cells expressing the mannosidase-Myc-HDEL from the strong constitutive GAP promoter.

Briefly, 0.5 g of wet weight yeast cells were lysed using 4×1 min vortexing with 4.5 g glass beads in 1 ml lysis-buffer (50 mM Tris-HCl pH 7.5 containing 0.6 M sorbitol, 10 mM β -mercaptoethanol and 5 mM MgCl₂). Between vortexing periods, the mixture was placed on ice for 5 min. The supernatant was collected and the glass beads were washed once with lysis-buffer, and the supernatant of this washing step was added to the first supernatant. This lysate was subjected to a differential centrifugation procedure. The P10000 pellet was solubilized in 0.5 ml of a 60% sucrose solution in lysis buffer. This solution was placed at the bottom of an Ultraclear ultracentrifuge tube (Beckman) of 14×89 mm. Subsequently, 1.5 ml each of sucrose solutions of 55, 50, 45, 42.5, 40, and 37.5% were carefully layered over each other. The tube was filled to the edge with 35% sucrose. Isopycnic sucrose gradient centrifugation was performed for 14 h at 180,000 g in a Beckman SW 41 rotor in a Beckman Model L8-70 preparative ultracentrifuge. After completion, 1 ml fractions were collected from the top and partially dialysed from excess sucrose, evaporated to dryness in a vacuum centrifuge. After redissolving the pellet in Laemmli buffer, the samples were subjected to SDS-PAGE in triplicate and the Western blots were treated with anti-HA, anti-Myc or anti-PDI ("PDI" for Protein Disulfide Isomerase), respectively.

The results illustrated almost exact cosedimentation of the MFManMycHDEL protein with the Protein Disulfide Isomerase marker protein (which is also targeted with a HDEL (SEQ ID NO: 1) signal sequence) (FIG. 5). In the same assay, the HA-tagged OCH1 was distributed over the whole gradient, with the highest abundance in fractions having a density lower than that of the fractions containing the mannosidase and the PDI. This result indicated that the mannosidase was targeted to the expected location (the ER-Golgi boundary) by the addition of an HDEL (SEQ ID NO: 1) signal. In contrast, the mannosidase without HDEL (SEQ ID NO: 1), expressed from inducible alcohol oxidase I promoter (which was of comparable strength as the GAP promoter), was secreted at a high level of about 20 mg/l.

Immunofluorescence Microscopy—

To confirm the correct targeting of the mannosidase-Myc-HDEL, an immunofluorescence microscopy experiment was performed.

Briefly, yeast cultures were grown to OD₆₀₀ in YPD (for pGAPZMFManMycHDEL) or in YMP following a YPGlycerol growth phase for pPICZBMFManMycHDEL. Formaldehyde was added to the yeast cultures to a final concentration of 4% and incubated for 10 min at room temperature. Cells were pelleted and resuspended in 50 mM potassium phosphate buffer pH 6.5 containing 1 mM MgCl₂ and 4% formaldehyde and incubated for 2 h at room temperature. After pelleting, the cells were resuspended to an OD₆₀₀=10 in 100 mM potassium phosphate buffer pH 7.5 containing 1 mM MgCl₂ and EDTA-free CompleteTM protease inhibitor cocktail (Boehringer Mannheim). To 100 μ l of cell suspension, 0.6 μ l of β -mercapto-ethanol and 200 of 20,000 U/ml Zymolyase 100T (ICN) were added, followed by a 25 minute incubation with gentle shaking. The cells were washed twice in the incubation buffer and added to poly-lysine coated cover slips (these are prepared using adhesive rings normally in use for reinforcing perforations in paper). Excess liquid was blotted with a cotton swab and the cells were allowed to dry at 20° C.

22

All blocking, antibody incubation and washing steps are performed in PBS containing 0.05% bovine serum albumin. Primary antibodies are used at 2 μ g/ μ l and secondary antibodies conjugated to fluorophores (Molecular probes) were used at 5 μ g/ μ l. The nucleus was stained with the nucleic acid stain HOECHST 33258. After fixation and cell wall permeabilization, the integrity of the yeast cell morphology was checked in phase contrast microscopy and after immunostaining, the slides were examined under a Zeiss Axiohot 10 fluororesense microscope equipped with a Kodak digital camera. Images were processed using Macprobe 4.0 software and prepared with Corel Photopaint 9.0.

The Golgi marker protein OCH1-HA gave the typical Golgi staining pattern described in the literature (speckle-like staining). Staining with the 9E10 monoclonal anti-Myc antibody, recognizing mannosidase-Myc-HDEL, gave a peri-nuclear staining pattern with some disparate staining in the cytoplasm, highly indicative for an ER targeting (FIG. 4).

Based on the foregoing experiments, it is concluded that 20 the *Trichoderma reesei* mannosidase-Myc-HDEL was targeted to the ER-Golgi boundary.

Example 2

Co-Expression of Mannosidase-HDEL with Recombinant Glycoproteins

Co-expression of Mannosidase-HDEL with the *Trypanosoma cruzi* trans-Sialidase

30 The cloning of a *Trypanosoma cruzi* trans-sialidase gene coding for an active trans-sialidase member without the C-terminal repeat domain has been described by Laroy et al. (*Protein Expression and Purification* 20: 389, 2000) which is incorporated herein by reference. The sequence of this *Trypanosoma cruzi* trans-sialidase gene is available through NCBI Genbank under the Accession No. AJ276679. For expression in *P. pastoris*, the entire gene was cloned in pHILD2 (Invitrogen, San Diego, Calif.), creating pHILD2-TS. To allow better secretion, pPIC9-TS was created in which 35 trans-sialidase was linked to the prepro secretion signal of the yeast α -mating factor. Plasmids pPIC9-TSE and pCAGGS-prepro-TSE were created where the epitope E-tag was added to the C-terminal of the trans-sialidase to allow easy detection and purification. The construction of pHILD2-TS, pPIC9- 40 TSE and pCAGGS-prepro-TSE has been described by Laroy et al. (2000), incorporated herein by reference. The vectors used in the construction were made available through for pCAGGS (No. LMBP 2453), Invitrogen, San Diego, Calif. for pHILD2 and pPIC9, and Pharmacia Biotech for 45 pCANTAB-5E.

Plasmid pPIC9-TSE was linearized with SstI and was transformed into *P. pastoris* GS115 (his4) strain by electroporation according to the manufacturer's instructions (Invitrogen). One of the transformants was further transformed 50 with plasmid pGAPZMFManHDEL, establishing a strain co-expressing Mannosidase-HDEL and the *Trypanosoma cruzi* trans-sialidase.

Fermentation and protein purification was according to the procedures described by Laroy et al. (2000).

Purified trans-sialidase was subject to carbohydrate analysis according to Callewaert et al., *Glycobiology* 11, 4, 275-281, 2001. Briefly, the glycoproteins were bound to the PVDF membrane in the wells of a 96-well plate, reduced, alkylated and submitted to peptide-N-glycosidase F deglycosylation. 55 The glycans were derivatised with 8-amino-1,3,6-pyrenetrisulfonic acid by reductive amination. Subsequently, the excess free label was removed using Sephadex G10-packed

23

spin columns and the glycans were analysed by electrophoresis on a 36 cm sequencing gel on an ABI 377A DNA-sequencer and detected using the built-in argon laser. Digests with 3 mU/ml purified *T. reesei* α -1,2-mannosidase (described by Maras et al., *J. Biotechnol.* 77, 255-63, 2000) were also performed in 20 mM sodium acetate pH=5.0. The glycans derived from 1 μ g of the purified recombinant glycoproteins were used as the substrate. 1 U of the α -1,2-mannosidase is defined as the amount of enzyme that releases 1 μ mol of mannose from baker's yeast mannan per minute at 37° C. and pH=5.0.

As can be seen in FIG. 6, panel B, the major N-glycan on trans-sialidase was Man₈GlcNAc₂ (Compare with panel F, representing an analysis of the N-glycans of bovine RNaseB. The one but last peak in this profile is Man₈GlcNAc₂, the first peak is Man₅GlcNAc₂). In vitro, this glycan was digestible to Man₅GlcNAc₂ with α -1,2-mannosidase (FIG. 6, panel C). In the N-glycan profile of the trans-sialidase co-expressed with mannosidase-HDEL, the major peak corresponded to Man₅GlcNAc₂ (FIG. 6, panel D).

Co-Expression of Mannosidase-HDEL with the Influenza a Virus Haemagglutinin

The Influenza A virus haemagglutinin was known to be glycosylated in *Pichia pastoris* with high-mannose N-glycans containing 9-12 mannose residues (Saelens et al. *Eur. J. Biochem.* 260: 166-175, 1999). The effect of a co-expressed mannosidase on the N-glycans of the haemagglutinin was assessed in an N-glycan profiling method described below. In addition, to compare the efficiency of the *Trichoderma* enzyme (having a temperature optimum of 60° C.) with a mammalian mannosidase having a temperature optimum of 37° C., the catalytic domain of the mouse mannosidase IB from a mouse cDNA-library was cloned and tagged with a HDEL signal by PCR amplification. This ORF was cloned after the prepro-signal sequence of the *S. cerevisiae* α -mating factor under the control of the GAP promoter. Expression of the mannosidase-HDEL transgenes on the mRNA level was confirmed by qualitative Northern blotting.

The haemagglutinin was expressed and purified from a non-mannosidase expressing control strain and from a strains co-expressing the *Trichoderma reesei* mannosidase-HDEL or the mouse mannosidase IB-HDEL according to the procedure described by Kulakosky et al. *Glycobiology* 8: 741-745 (1998). The purified haemagglutin was subjected to PNGase F digestion as described by Saelens et al. *Eur. J. Biochem.* 260: 166-175, 1999. The proteins and glycans were precipitated with 3 volumes of ice-cold acetone and the glycans were extracted from the pellet with 60% methanol. Following vacuum evaporation, the glycans were labeled with 8-amino-1,3,6 pyrenetrisulfonic acid by adding 1 μ l of a 1:1 mixture of 20 mM APTS in 1.2M citric acid and 1M N_aCNBH₃ in DMSO and incubating for 16 h at 37° C. at the bottom of a 250 μ l PCR-tube. The reaction was stopped by the addition of 10 μ l deionized water and the mixture was loaded on a 1.2 cm Sephadex G10 bed packed to dryness in a microspin-column by centrifugation in a swinging bucket rotor, which provided for a flat resin surface. After loading, 50 μ l deionised water was carefully added to the resin bed and the spin column was briefly centrifuged for 5 seconds at 750 g in a tabletop centrifuge. This elution process was repeated twice and all the eluates were pooled and evaporated to dryness in a Speedvac vacuum centrifuge (Savant). The labeled glycans were reconstituted in 1.5 μ l gel loading buffer containing 50% formamide and 0.5 μ l Genescan 500™, labeled with rhodamine (Perkin Elmer Bioscience), serving as an internal reference standard. This mixture was loaded on a DNA-sequencing gel containing 10% of a 19:1 mixture of acrylamide:bisacryla-

24

mide (Biorad, Hercules, Calif., USA) and made up in the standard DNA-sequencing buffer (89 mM Tris, 89 mM borate, 2.2 mM EDTA). Polymerization of the gel was catalyzed by the addition of 200 μ l 10% ammoniumpersulfate solution in water and 20 μ l TEMED. The gel was of the standard 36 cm well-to-read length and was run on an Applied Biosystems Model 373A DNA-sequencing apparatus. Pre-running of the gel was done at 1000 V for 15 min. and after loading, the gel was electrophoresed for 8 h at 1250 V without heating. This methodology gives a limit of detection of 10 fmol per peak. The data were analysed with Genescan 3.0 software.

As shown in FIG. 7, the *Trichoderma reesei* α -1,2-mannosidase provided the most complete reduction in the number of α -1,2-mannoses present on the N-glycans. The N-glycan processing by mouse mannosidase IB-HDEL was less efficient than by the *Trichoderma reesei* α -1,2-mannosidase.

Despite the efficient removal of α -1,2-mannoses from the N-glycans of haemagglutinin, no Man₅GlcNAc₂ was obtained. Even after digestion of the N-glycans with 3 mU of purified *Trichoderma reesei* α -1,2-mannosidase, only Man₆GlcNAc₂ was obtained as the smallest sugar chain. These results indicated that the remaining residues were possibly α -1,6-linked mannoses, originating from the initiating OCH1 α -1,6-mannosyltransferase enzymatic activities. OCH1 was observed to be localized to very early part of the Golgi apparatus and could act on the N-glycans of haemagglutinin before complete digestion of the Man₈GlcNAc₂ precursor to Man₅GlcNAc₂ by the mannosidases-HDEL. Thus, for proteins whose glycans are efficiently modified by the α -1,6-mannosyltransferase, an inactivation of the OCH1 gene coding for the transferase would be desirable in order to obtain proteins with Man₅GlcNAc₂.

Example 3

Inactivation of the *Pichia* Och1 Gene

A *Pichia pastoris* sequence was found in the GenBank under Accession No. E12456 and was described in Japanese Patent Application No. 07145005, incorporated herein by reference. This sequence shows all typical features of an α -1,6-mannosyltransferase and is most homologous to the *S. cerevisiae* OCH1, thus referred to herein as the *Pichia pastoris* Och1 gene.

First, the full ORF of the *Pichia pastoris* Och1 gene was PCR cloned in pUC18 to obtain plasmid pUC18pOch1. pUC18pOch1 was cut with HindIII, blunt-ended with T4 polymerase, then cut with XbaI, releasing a fragment containing the 5' part of the *Pichia pastoris* Och1 gene. This fragment was ligated into the vector pBLURA IX (available from the Keck Graduate Institute, Dr. James Cregg, which had been cut with Eco RI, blunt-ended with T4 polymerase, and then cut with Nhe I. This ligation generated pBLURA5'PpPCH1, as shown in FIG. 8.

Disruption of this *Pichia* OCH1 gene in the *Pichia* genome was achieved by single homologous recombination using pBLURA5'PpOCH1, as illustrated in FIG. 9. As a result of the single homologous recombination, the Och1 gene on the *Pichia* chromosome was replaced with two Och1 sequences: one consisted only about the first one third of the full Och1 ORF, the other had a full Och1 ORF without a Och1 promoter. Single homologous recombination was achieved as follows. Cells of the *Pichia* strain yG4 were transformed by electroporation with pBLURA5'PpOCH1 which had been linearized with the single cutter Bst BI. About 500 transformants were obtained on minimal medium containing 1M sorbitol,

25

biotin, arginine, adenine and histidine and incubation at 27° C. Thirty-two of these transformants were picked and re-selected under the same conditions. Twelve clones were further analyzed for correct genomic integration of the cassette by PCR. Seven of the twelve URA prototrophic clones contained the cassette in the correct locus.

One of the Och1-inactivated clones was also further transformed with pGAPZMFManHDEL to produce "supertransformants". Both the Och1-inactivated clone and three super-transformants also expressing the ManHDEL were evaluated in cell wall glycan analysis as follows. Yeast cells were grown in 10 ml YPD to an OD₆₀₀=2 and mannoproteins were prepared by autoclaving the yeast cells in 20 mM sodium citrate buffer pH 7 for 90 min at 120° C. and recovery of the supernatant after centrifugation. Proteins were precipitated from this supernatant with 3 volumes of cold methanol. The protein preparation obtained in this way was used for N-glycan analysis using DSA-FACE as described by Callewaert et al. (2001) *Glycobiology* 11, 275-281. As shown in FIG. 10, there was an increased amount of Man₈GlcNAc₂ glycan in the Och1-inactivated clone as compared to parent strain yGC4, indicative of a reduced activity of the Och1 enzyme. In all three super-transformants which also expressed the HDEL (SEQ ID NO: 1)-tagged α-1,2 mannosidase, the production of Man₅GlcNAc₂ was observed. Furthermore, upon digestion of the same glycan mixtures with 3 mU/ml purified recombinant *Trichoderma reesei* α-1,2-mannosidase, more Man₅GlcNAc₂ was formed in the strain transformed with pBLURA5'PpOCH1 than in the parent strain (FIG. 11, compare panel 2 and 3).

These results confirmed that the lack of a production of Man₅ glycans on recombinantly produced proteins such as haemagglutinin from cells expressing α-1,2-mannosidase were due to the activity of the Och1 protein. These results further indicate that the production of glycoproteins with Man₅ glycans could be facilitated by the inactivation of the Och1 gene.

Example 4

Expression of Glucosidase II in *Pichia pastoris*

4.1 Amplification of the GLSII Alpha Subunit ORF from *S. cerevisiae*.

Genomic DNA was prepared from the *S. cerevisiae* strain INVS (α, leu2-3, 112 h is 3Δ1, trp1-289, ura3-52), using the Nucleon kit (Amersham). A touch-down PCR reaction was performed using this genomic DNA as template and the LA TaKaRa polymerase (ImTec Diagnostics). The sequence of the PCR primers was based on the known sequence of the *S. cerevisiae* GLSII ORF:

Sense primer:

(SEQ ID NO: 12)
5' CCG CTC GAG ATG GTC CTT TTG AAA TGG CTC 3'
Xho I

Antisense primer:

(SEQ ID NO: 13)
5' CCG GGC CCA AAA ATA ACT TCC CAA TCT TCA G 3'
Apa I

4.2 Cloning of the *S. cerevisiae* Glucosidase II ORF into *Pichia pastoris* Expression Vectors.

Construction of the Glucosidase II Expression Vectors—

The PCR fragment was digested with Xho I/Apa I and ligated into the pGAPZA vector (Invitrogen), thereby placing the ORF under the transcriptional control of the GAP pro-

26

moter. Using this strategy, the myc and the His6 tag were placed in frame to the C-terminus of Glucosidase II, creating pGAPZAGLSII. The complete ORF of pGAPZAGLSII was then sequenced to ensure that no mutations were generated in the PCR reaction. The sequence of the vector pGAPZAGLSII was set forth in SEQ ID NO: 18. The GLSII ORF from the pGAPZAGLSII vector was cloned into vector pPICZA (Invitrogen) to create pPICZAGLSII, thereby placing the ORF under the transcriptional control of the AOX1 promoter. The GLSII ORF from the pGAPZAGLSII vector was cloned into vector pAOX2ZA, thereby placing the ORF under the transcriptional control of the AOX2 promoter. This vector was created by replacing the multi cloning site of vector pAOX2ZB with the multi cloning site of pPICZA. Vector pAOX2ZB was generated by replacing the AOX1 promotor of pPICZB by the AOX2 promotor region of the AOX2 gene (Martinet et al., Biotechnology Letters 21). The AOX2 promotor region was generated by PCR on *Pichia* genomic DNA with the sense primer 5'GACGAGATCTTTTCAGAC-CATATGACGG 3' (SEQ ID NO: 26) and the antisense primer 5'GCGGAATTCTTCAGTTGATTTGT 3' (SEQ ID NO: 27). The GLSII ORF from the pGAPZGLSII vector was cloned into vector pYPT1ZA to create pYPT1-ZAGLSII, thereby placing the ORF under the transcriptional control of the YPT1 promoter. Vector pYPTZA was created by replacing the AOX1 promotor of pPICZA by the YPT1 promoter present on the plasmid pIB3 (GenBank accession number AF027960)(Sears et al., Yeast 14, pg 783-790, 1998). All constructs contain the phleomycin resistance gene. The resulting final expression vectors (pGAPZAGLSII, pAOX2ZAGLSII, pPICZAGLSII and pYPT1ZAGLSII) are depicted in FIGS. 12-15.

Similar expression vectors were constructed, carrying the Ampicillin resistance marker and the *Pichia* ADE1 selection marker. In principle, the Zeocin resistance expression cassette of the plasmids pAOX2ZAGLSII, pGAPZAGLSII and pYPT1ZAGLSII was replaced by the Ampicillin and *Pichia* ADE1 cassette of the vector pBLADE IX (Cregg, J. M.) to result in the vectors pAOX2ADE1glsII, pGAPADE1glsII and pYPT1ADE1glsII. Vector pPICADE1glsII was obtained by inserting the glucosidase II open reading frame into the multiple cloning site of the vector pBLADE IX (Cregg, J. M.). The resulting final expression vectors (pGAPADE1glsII, pAOX2ADE1glsII, pPICADE1glsII and pYPT1ADE1glsII) are depicted in FIGS. 16-20.

Adding the ER Retention Tag HDEL to Glucosidase II Expression Vectors—

The following primers were used to generate an HDEL-containing PCR fragment:

Primer 1:

(SEQ ID NO: 28)
5' GCG GGT CGA C/CA C/GA C/GA A/CT G/TG A/GT TTT
Sal I H D E L stop
AGC CTT AGA CAT GAC 3'

Primer 2:

(SEQ ID NO: 29)
5' CAG GAG CAAA GCT CGT ACG AG 3'
SpI I

PCR was performed on pGAPZAGLSII with Taq pol., at 60° C. The PCR fragment of 225 bp was cut with Sal I/Spl I and ligated into the Sal I/Spl I opened pGAPZAGLSII vector, creating plasmid pGAPZAglsIIHDEL. The sequence of plasmid pGAPZAglsIIHDEL is set forth in SEQ ID NO: 24. The

27

construction strategy and the resulting final expression vectors (pGAPZAglsIIHDEL and pGAPADE1glsIIHDEL) are depicted in FIGS. 20-21.

4.3 Transformation of a *Pichia pastoris* Strain.

Transformation was performed using the conventional electroporation techniques, as described by Invitrogen. Cells of the *Pichia pastoris* strain PPY12-OH were transformed with pGAPZGLSII which had been cut with the single cutter Avr II. Transformants were selected based on their resistance to zeocin.

Genomic Analysis of the Transformants—

Genomic DNA was prepared from some zeocin resistant *Pichia* transformants. A PCR reaction was performed on the genomic DNA in order to determine whether or not the glucosidase II gene was integrated into the yeast genome. PCR was performed using Taq DNA polymerase (Boehringer) (2.5 mM MgCl₂, 55° C. for annealing). The primers were the same as the ones we used for the amplification of the ORF on *S. cerevisiae* genomic DNA. pGAPZAGLSII transformants were confirmed by the presence of a specific PCR product indicative of the glucosidase II ORF.

4.4 Expression and Secretion of the *S. Cerevisiae* Glucosidase II Alpha Subunit in *Pichia pastoris*

Analysis at the Transcriptional Level—

RNA was prepared from the transformants which scored positive after the genomic analysis. RNA was prepared using acid phenol. From each sample, 15 µg of RNA was loaded on a formaldehyde agarose gel. After electrophoresis the RNA was blotted on a Hybond N membrane. The membrane was hybridizing using a radioactive probe, which consists of a 344 bp glucosidase II specific fragment, corresponding to the 3' region of the glucosidase II ORF. No signals were detected with non-transformed control strains, whereas clear signals were observed with transformants.

Analysis at the Protein Level Using a Double Membrane Assay—

A nitrocellulose membrane was placed on a buffered dextrose medium (BMDY). On top of that nitrocellulose membrane, a cellulose acetate membrane was placed. *Pichia* transformants of pGAPZAGLSII were streaked on the cellulose acetate and grown for a few days. The yeast cells remained on the cellulose acetate, while the secreted proteins crossed this membrane. As such the secreted protein was captured onto the nitrocellulose membrane. After a few days the cellulose acetate, containing the yeast colonies, was removed. The nitrocellulose membrane was analyzed for the presence of glucosidase II using anti-myc antibody. Most of the transformants gave a clear signal as compared to a faint, hardly visible signal with the WT, non-transformed strain.

Extracellular Expression—

PPY12-OH transformants of the construct pGAPZAGLSII (mychis6) (strains 12, 14 and 18) and transformants of the construct pGAPZAGLSII(myc)HDEL (strains H1, H2 and H3) were grown for 2 days on 2×10 ml BMDY medium. These 6 transformants earlier scored positive both on the genomic level (PCR on gDNA) and on the RNA level (Northern blot). The culture medium was collected by centrifugation and concentrated with Vivaspin columns to about 1 ml. Proteins from this concentrate were precipitated with TCA, resuspended in Laemmli buffer and loaded for SDS-PAGE analysis. Proteins were blotted to nitrocellulose membrane. The blot was incubated overnight with anti-myc Ab. The secondary Ab was linked to peroxidase. Using the Renaissance luminiscence detection kit (NEN) and a light sensitive film (Kodak), a strong band at about 110 kDa was observed for the transformants 12, 14 and 18, indicating that GLSII was expressed and secreted from these transformants. No signal

28

was obtained for the transformants H1-3, which indicate that the HDEL (SEQ ID NO: 1) tag, which was added C-terminally to the GLSII ORF, resulted in an ER localization of the protein, preventing GLSII to be secreted into the growth medium.

Intracellular Expression—

The 6 transformants and the WT strain were grown for 2 days in 500 ml BMDY. The cells were collected by centrifugation, washed, resuspended into a minimal volume (50 mM Tris.HCl pH 7.5, 5% glycerol) and broken using glass beads.

The cell debris was removed through several centrifugation steps (low speed centrifugation (2000-3000 g)). Membranes were obtained from the supernatant through ultracentrifugation. The pellets were resuspended in Laemmli buffer and loaded for SDS-PAGE analysis. The proteins were blotted on a nitrocellulose membrane. The intracellular GLSII expression was checked using anti-myc Ab and peroxidase conjugated secondary Ab. Following the luminescence detection, a band at about 110 kDa was observed with the GLSIIHDEL transformants (H1 and H3, faint signal for H2), but not with the WT and GLSII expression strains. These results clearly indicate the intracellular presence of the recombinant GLSII when expressed with a C-terminal HDEL (SEQ ID NO: 1) tag. No GLSII was detected intracellularly when this tag was not present.

4.5 Purification and Activity Assays of the Recombinant Glucosidase II Alpha Subunit

A GLSII assay was set up as follows and was tested using a commercially available yeast alpha-glucosidase (Sigma) as a positive control.

Composition: 70 µl 80 mM phosphate-citrate buffer pH 6.8, 7 µl 250 mM mannose, 3.5 µl 250 mM 2-deoxy-D-glucose, 0.8 µl 4-MeUmbelliferyl-alpha-D-glucopyranoside (1 µM). Three assays were performed: one with 1 unit commercial enzyme, one without the enzyme and one with the enzyme but without the substrate. The assay mixture was incubated overnight at 30° C. When illuminated with UV, only the reaction mixture with both the enzyme and the substrate showed fluorescence (FIG. 22). This indicates that the assay was very specific in detecting the activity of the alpha-glucosidase.

WT PPY12-OH, strain 18 and strain H3 were grown during 2 days in 2×10 ml growth medium. Cells were spun down and medium was adjusted to 300 mM NaCl and 10 mM imidazol and concentrated with Vivaspin columns to 0.5-1 ml. Medium was loaded onto a Ni-NTA spin column (Qiagen) and the purification was performed according to the manufactures recommendations. Protein was eluted from the column in 2×100 µl elution buffer (50 mM Na₂PO₄, 300 mM NaCl, 250 mM imidazol pH 8.0). From each eluate, 20 µl was assayed for its glucosidase II activity. 0.33 units of the commercial enzyme diluted in 20 µl of the elution buffer was used as a positive control. Fluorescence was observed with the positive control and the elute of strain 18, the strain which secreted the enzyme into the growth medium. These results indicate that the recombinant *S. cerevisiae* GLSII alpha subunit, secreted by *Pichia pastoris*, was a functionally active enzyme. The activity was not seen in the WT (untransformed) strain, nor in strain H3 as the GLSII was expressed intracellularly (FIG. 23). These results also indicate that the beta subunit is not necessary for the functionality of the alpha part of the protein.

Example 5

Creating *Pichia* Strains Expressing Both Glucosidase II and Mannosidase

Strain GS 115 was transformed with pGAPZGLSII and pGAPZglsIIHDEL. Transformants were selected on YPD-Szeo.

29

Strain yGC4 was transformed with the following constructs, respectively:

- (1) pGAPADEglsII and pGAPADEglsIIHDEL, selection on synthetic sorbitol medium without adenine;
- (2) pGAPZMFMManHDEL; selection on YPDsZeo; and
- (3) pGAPZMFMManHDEL/pGAPADEglsIIHDEL: selection on synthetic sorbitol medium without adenine and with zeocin.

30

Strain yGC4 with OCH1 knock-in and expressing MFmannosidaseHDEL was transformed with pGAPADEglsII and pGAPADEglsIIHDEL. Selection of transformants was done on synthetic sorbitol medium without adenine and uracil.

For all transformations, colonies were obtained. Transformants with the expression vector(s) integrated into the genome, determined by PCR, were obtained. Expression of GLSII from some of these transformants was observed.

SEQUENCE LISTING

```
<160> NUMBER OF SEQ ID NOS: 29
<210> SEQ ID NO 1
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:synthetic
peptide representing the ER-retention signal.
```

```
<400> SEQUENCE: 1
```

```
His Asp Glu Leu
1
```

```
<210> SEQ ID NO 2
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer
```

```
<400> SEQUENCE: 2
```

```
gactggttcc aattgacaag c
```

21

```
<210> SEQ ID NO 3
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer
```

```
<400> SEQUENCE: 3
```

```
agtcttagatt acaactcgtc gtgagcaagg tggccgcccc gtcg
```

44

```
<210> SEQ ID NO 4
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer
```

```
<400> SEQUENCE: 4
```

```
ccattgagga cgcgtccgc gcc
```

23

```
<210> SEQ ID NO 5
<211> LENGTH: 85
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer
```

```
<400> SEQUENCE: 5
```

```
gtatctagat tacaactcgt cgtgcagatc ctcttctgag atgagtttt gttcagcaag
```

60

```
gtggccgccc cgtcgtatc atgaa
```

85

```
<210> SEQ ID NO 6
```

-continued

<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 6

aactcgagat ggactcttca aaacacaaaac gc

32

<210> SEQ ID NO 7
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 7

ttgcggccgc ttacaactcg tcgtgtcggg caggaggatt acctga

46

<210> SEQ ID NO 8
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 8

ccattgagga cgcatgccgc gcc

23

<210> SEQ ID NO 9
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 9

gcaaatggca ttctgacatc ct

22

<210> SEQ ID NO 10
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 10

gtcccttattt caatcaatttg aa

22

<210> SEQ ID NO 11
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 11

gactgggtcc aattgacaag c

21

<210> SEQ ID NO 12
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 12

-continued

ccgctcgaga tggtcctttt gaaaatggctc 30

<210> SEQ ID NO 13
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 13

cggggcccaa aaataacttc ccaatcttca g 31

<210> SEQ ID NO 14
<211> LENGTH: 1785
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:The ORF sequence of the MFManHDEL fusion in pGAPZMFManHDEL.

<400> SEQUENCE: 14

at gagat tt cttcaat tt tactgctgtt ttattcgag catcctccgc attagotgct	60
cc agtcaaca ctacaacaga agatgaa acg gcacaaat tc cggctgaagc tgtcatcggt	120
tactcagatt tagaagg gga tt tcgatgtt gctgtttgc cattttccaa cagcacaat	180
aacgggttat tg ttataaa tactactatt gccagcatg ctgctaaga agaagg gta	240
tctctcgaga aa aga gagggc tgaagctgaa tt cgccacaa aacgtggatc tcccacccct	300
acgagg ggcgg cagcagtcaa ggccgcattc cagacgtcgt ggaacgctta ccaccat	360
gccttcccc atgacgaccc ctacccggc agcaacagct ttgatgatga gagaacggc	420
tggggctegt cggcaatcga tggcttgac acggctatcc tcatggggta tgccgacatt	480
gtgaacacga tcctttagta tgtaccgag atcaacttca ccacgactgc gtttgcac	540
caaggatcct ccgtgttgcgac gaccaacatt cggtaccctcg gtggccgtct ttctgcctat	600
gacctgttgc gaggtccctt cagctccctg cgcacaa acc agaccctgtt aaacagcctt	660
ctgaggcagg ctcaaacact ggccaaacggc ctcaagggtt cgttccacac tcccacgg	720
gtcccggaec ctaccgtt cttcaaccctt actgtccggaa gaagtgggtc atctagcaac	780
aacgtcgctg aaatttggaa cctgggtcgtc gagtggacac ggtttagcga cctgacgg	840
aaccccgagt atgcccagct tgccgagaag ggccgactgt atctcctgaa tccaaagg	900
agcccgagg catggccctgg cctgatttgc acgtttgtca gcacgagca cggta	960
caggatagca gcccggactg gtccggcctc atggacagct tctacgatca cctgatcaag	1020
atgtacccgt acgaccggcgt tgctttgcactacaagg atcgctgggt ctttgggt	1080
gactcgacca ttgggcatct cggctctc acgtcgacgc gcaaggactt gacccctt	1140
tcttcgtaca acggacagtc tacgtcgcca aactcaggac atttggccag ttttggcg	1200
ggcaacttca tcttggagg cattctcctg aacgagcaaa agtacattga ctttggatc	1260
aagcttgcac gctgtactt tggcacgtac acccagacgg cttctggaa cggccocgaa	1320
ggcttcgcgt ggggtggacag cgtgacggc gcccggcgt cggccgcctc gtcccgatcc	1380
gggttctact cgtcgccagg attctgggtt acggcaccgtt attacatctt gcccggag	1440
acgctggaga gcttgtacta cgcataccgc gtcacggcgc actccaagtg gcaggac	1500
gcgtggaaag cgttgagtgc cattgaggac gcatgcccgc cggcagcgc gtactcg	1560

-continued

atcaacgacg tgacgcaggc caacggcggtt ggtgcctctg acgatatgga gagcttcgtt	1620
tttgcgcagg cgctcaagta tgcgtacctt atctttgcggg aggagtcggg tgtgcagggtt	1680
caggccacccg gcgggaacaa atttgtctt aacacggagg cgcacccctt tagcatccgt	1740
tcatcatcac gacggggggcc ccaccccttgct cacgacgagt tgtaa	1785
<210> SEQ ID NO 15	
<211> LENGTH: 2016	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence: The ORF sequence of the MFmManHDEL fusion in pGAPZMFMmManHDEL.	
<400> SEQUENCE: 15	
atgagatttc cttcaatttt tactgctgtt ttatcgtag catcctccgc attagctgtt	60
ccagtcaaca ctacaacaga agatgaaacg gcacaaaatc cggctgaagc tgtcatcggt	120
tactcagatt tagaaggggtt tttcgatgtt gctgtttgc cattttccaa cagcacaat	180
aacgggttat tgtttataaa tactactatt gccagcatg ctgctaaaga agaagggtt	240
tctctcgaga tggactcttc aaaacacaaa cgctttgatc tgggctttaga agatgtgtt	300
attcctcactg tagatgccgg caaaggagct aaaaaccccg gcgtcttcctt gatccatgg	360
cccgacgaac acagacacag ggaagaagaa gagcgtctga gaaataagat tagagctgac	420
catgagaaag ccctggaaaga agcaaaagaa aaattaagaa agtcaagaga ggaaatccgt	480
gcagaaattc agacagagaa aaacaaagta gccaagcaa tgaagacaaa agagaccagg	540
gtactgccgc ctgtccctgt cccacaacgt gttaggggtca gtgggtgggaa tccagaagac	600
atggagatca agaagaaaag agacaaaattt aaagagatga tgaacacatgc ctgggataat	660
tacagaacat acggatgggg acataatgaa ctaaggccta ttgcaaggaa aggccattcc	720
actaacatata tcggaagctc acagatgggtt gccaccatag tggatgtttt ggataccctt	780
tatatacatgg ggcttcatga tgaattcatg gatggggaaa gatggattga agaaaacctt	840
gatttcagtg tgaattcaga agtgtctgtc ttgaagtta acattcgctt tattggaggg	900
ctccctcgctg catattacct gtcaggagag gaaatattca agactaaagc agtgcagtt	960
gctgagaaac tccttcctgc cttaacaca cctactggga ttcccctggc aatggtaac	1020
ctgaaaagtg gagtaggtcg aaactggggc tggcgtctg caggcagcag catcctggct	1080
gagttccggca ccctgcacat ggagtttgcg cacctcagct acttgaccgg tgacttgact	1140
tactataata aggtcatgca cattcggaaa ctactgcaga aaatggaaacg cccaaatgg	1200
ctttatccaa attattnaa cccaaagaaca gggcgctggg gtcagatca cacaatcgat	1260
ggtgtctgg gagatagttt ttatgaatac ttactgaaag catggctgac gtcagataaa	1320
acagaccacg aggcaagaag gatgtatgac gatgctgtt aggctataga aaaacatctt	1380
atataagaagt cccgaggagg tctggttttt attggagaat ggaagaatgg acacttggaa	1440
aggaagatgg ggcacttggc ctgctttgtc gggggaaatgc ttgccttgg agcagatgg	1500
tccagaaagg ataaagctgg ccactactta gaacttagggg cagaaatgc acgaacatgt	1560
catgagtcat atgacagaac tgcattgaaa ctaggtccgg agtcattcaa gtttgcgtt	1620
gcagtggaaag ccgtggctgtt gggcgggtt gaaaaggattt acatcctcg tccagaagta	1680
attgaaacctt attggatctt atggcgattt acccacgacc caagatacag gcagtggggc	1740
tggaaagcag cactggctat tgagaagtcg tgccgggtca ggggtgggtt ttctgggtc	1800

-continued

aaggatgtat acgccccgac ccctgtgcac gacgacgtgc agcagagactt ttctttgtct	1860
gaaacattaa aatacttgta cctgtgttc tctggcgatg accttctacc tttagaccac	1920
tgggtgttta acacagaggc gcaccctctg ccgggtgtgc gcttagccaa cagcaactt	1980
tcaggttatac ctgtgttccg acacgacgag ttgtaa	2016

<210> SEQ ID NO 16
<211> LENGTH: 6757
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pAOX2ZAGLSII.

<400> SEQUENCE: 16	
catggccaag ttgaccagtgc cggttccgggt gtcaccgcg cgacgtgc ccggagcggt	60
cgggttctgg accgaccggc tcgggttctc ccgggacttc gtggaggacg acttcgcgg	120
tgtggtcgg gacgacgtga ccctgttcat cagcgcggc caggaccagg tggtgcgg	180
caacaccctg gcctgggtgt gggtgcgcgg cctggacgag ctgtacggcc agtggtcgg	240
ggtcgtgtcc acgaacctcc gggacgcctc cgggcggcc atgaccgaga tcggcgagca	300
gccgtggggg cgggagttcg ccctgcgcga cccggccggc aactgcgtgc acttcgtggc	360
cgaggagcag gactgacacg tccgacggcg gcccacgggt cccaggcctc ggagatccgt	420
ccccctttt ctttgtcat atcatgtat tagttatgtc acgcttacat tcacgcctc	480
cccccacatc cgctctaacc gaaaaggaaag gagttagaca acctgaagtc taggtcccta	540
tttattttt tatagttatg ttagtattaa gaacgttatt tatatttcaa atttttttt	600
tttttctgtat cagacgcgtg tacgcgtatc acattataact gaaaacccctt cttgagaagg	660
ttttgggacg ctcgaaggct ttaatttgc a gctggagac caacatgtga gcaaaaggcc	720
agcaaaaggc caggaaccgt aaaaaggccg cgttgcgtgc gttttccat aggctccgccc	780
ccccgtacga gcatcacaaa aatcgacgtc caagtcagag gtggcgaaac ccgcacaggac	840
tataaagata ccaggcggtt cccctggaa gtcctctgt gcgctctctt gttccgaccc	900
tgcgcgttac cggataacctg tccgccttc tcccttcggg aagcgtggcg ctttctcaat	960
gctcacgtc taggtatctc agttcggtgt aggtcggtcg ctccaaatgtc ggctgtgtgc	1020
acgaaccccc cgttcagccc gaccgcgtcg ctttccgg taactatgtc ttgagtc	1080
acccggtaag acacgactta tcgcactgg cagcagccac tggtaacagg attacgag	1140
cgaggtatgt aggcgggtgtc acagagttct tgaagtgggt gcctaactac ggctacacta	1200
gaaggacagt atttggatc tgcgtctgc tgaaggccat taccttcggg aaaagatgt	1260
gtagcttttgc atccggcaaa caaaccaccc ctggtagccg tggttttttt gtttgc	1320
agcagattac ggcgcgaaaa aaaggatctc aagaagatcc tttgatctt tctacgggt	1380
ctgacgtca gtggAACGAA aactcacgtt aagggtttt ggtcatgaga tcagatctt	1440
ttttcagacc atatgaccgg tccatcttc acggggggat tatctatgtt ttgacccat	1500
tcttgattct ttatgtattt aaatcacatc tacgttattt attacttact ggttatttac	1560
ttagcgcctt ttctgaaaaa catttactaa aaatcataca tcggcactct caaacacgac	1620
agattgtat caagaaggcag agacaatcac cactaagggtt gcacatttgc ggcaggc	1680
tcctaataga ggttcgatac ttatgtat aatacgacat attgtctac ctctgaatgt	1740
gtcaataactc tctcgatcc cgttcgtca gctaaaaata taacacttcg agtaagatac	1800

-continued

gcccaattga aggctacgag ataccagact atcactagta gaactttgac atctgtaaa	1860
gcagatcaaataccat tccagaatca attacccccc tttagctgtt cgaaggcatg	1920
aaaaagctac atgaaaatcc ccatccttga agtttgta gcttaaagga ctccattcc	1980
taaaatttca agcagtccctc tcaactaaat tttttccat tcctctgcac ccagccctct	2040
tcatcaaccg tccagccccc tcaaaagtcc aatgttaagta gcctgcaaat tcaggttaca	2100
acccctcaat ttccatcca agggcgatcc ttacaaagtt aatatcgaa acgagagact	2160
aaggagtc a tcatcaccc ccaacgatgg tgaaaaactt taagcataga ttgtatggagg	2220
gtgtatggca cttggcggtt gcatttagt ttgaaactat ggggtaataac atcacatccg	2280
gaactgatcc gactccgaga tcatatgcaaa agcacgtgat gtaccccgta aactgctcg	2340
attatcgttt caattcatcg tcttaaacag tacaagaaac ttatttcatg ggtcatttgg	2400
ctctgatgag gggcacattt ccccaatgtat ttttggaa agaaagccgt aagaggacag	2460
ttaagcgaaa gagacaagac aacgaacagc aaaagtgaca gctgtcagct accttagtgg	2520
cagttggag ttccaaattt gttgggtttt aatttttacc catgttgagt tgccttgct	2580
tctccttgca aacaatgcaaa gttgataaga catcacccccc caagataggc tattttgtc	2640
gcataaattt ttgtctcgga gtggaaaaccctt ttttatgtt aacagattac agaaggtcc	2700
tacccttcac cggtttagat ggggagaaaa ttaagcgatg aggagacat tattttgtata	2760
aaagaagcaa cccaaatccc ttattgtcct ttctgatca gcatcaaaga atattgtctt	2820
aaaacgggtt ttaactaca ttgttcttac acattgcaaa cctcttcctt ctatttcgga	2880
tcaactgtat tgactacatt gatctttttt aacgaagttt acgacttact aaatccccac	2940
aaacaaatca actgagaaaa gaattcacgt ggcccgccg gccgtctcg atcggtacct	3000
cgagatggtc ctttggaaat ggctcgatg ccaattggc ttctttaccg cttttcgca	3060
tgcgttacc gactatctat taaaagatgtt tgcgcaatctt gggttttgcc atagaaacag	3120
gtttagtgc aaaaatattt cccaaatctca tcaactgtat tacaatgtgg acgcccgtc	3180
tattgcacac gatccttttag agaatgtgtt tcatgttacc ataattaaaa ctataccaag	3240
attggagggc gatgatatacg cccattcttctt ctctctttt tacaggatca	3300
ctcagtaagg ttcaactataa atgagaaaaga gagaatgcca accaacacga gcggttgg	3360
gatctcttca caacgggttca atgagacctg gaagtacgca ttgcacaaga aatttcaaga	3420
ggaggcgaac aggaccagta ttccacaattt ccacttcctt aagcaaaaac aaactgtgaa	3480
ctcattctgg tcgaaaatat ttccat ttcactttca aactccactg cagacacatt	3540
tcatcttcga aacgggtatg tatccgtaga aatctttgtt gaacccccc aattgaaatg	3600
ttactggcaa aatgcgttca aacttattgtt aacgcggaa aatttcgttca acatttgcac	3660
tcatagaact aagcaggaaa acttcgcaca cgtgctgcca gaagaaacaa ctttcaacat	3720
gtttaaggac aatttcttgcatttca tgggttcttac taatgttctac ggtataccgg aacatgcgt	3780
tgcgttagat ttctcttca tgggttcttac taatgttctac ggtataccgg aacatgcgt	3840
gtcgctttagt ctgttgcata cttcaggatggaa aagggaaaccctt tacaggatgg tcaacgttgc	3900
tgtcttttagt tacaacatcg gtaccagccca accaatgtac ggttcgttcc cattcatgtt	3960
ttcatcttcg tccacatcta tctttttgggtt caatgcgtt gacacttgggg tagacataaa	4020
gtatgacacc agtaaaaataaa aacgcgttca tcatgttgcata cccggaaaatgttgc	4080
tgtatgttgc tccctggggc cagatattcc aactatcattt gacaaatgttgc	4140

-continued

tggtagaccc	tttttacccgc	ccatttcctc	tatagggtac	catcaatgt	gatgaaatta	4200
taatgatgag	atggacgttc	tcacagtgg	ctctcagatg	gatgctata	tgatttccta	4260
cgattttatt	tggttggact	tggagtatac	gaacgacaaa	aatatttt	cttggaaagca	4320
gcactccccc	cccaatccaa	aaaggctgtt	atccaaat	aaaaagtgg	gtagaaatct	4380
tgtcgacta	atcgatcctc	atttaaagaa	agattatgaa	atcagtgaca	gggttaattaa	4440
tgaaaatgt	gcagtcaagg	atcacaatgg	aatgactat	gtaggtcatt	gctggcagg	4500
taattctata	tggattgata	ccataagcaa	atatggccaa	aagatttgg	agtccctttt	4560
cgaacggttt	atggatctgc	cggctgattt	aactaattt	ttcatttgg	atgatatgaa	4620
cgagccttcg	atttcgatg	gcccagagac	cacagctcca	aaagatttga	ttcacgacaa	4680
ttacattgag	gaaagatccg	tccataacat	atatggctca	tcagtgcat	aagctactta	4740
cgacgcaata	aaatcgattt	attcaccatc	cgataagcgt	ccttccttc	taacaaggc	4800
ttttttgcc	ggctctcaac	gtactgctgc	cacatggact	ggtgacaatg	tggccaattt	4860
ggattactta	aagatttca	ttccttatgg	tctgtcaac	aacattgtcg	gtatgccatt	4920
tataggagcc	gacatagctg	getttgtga	ggatcctaca	cctgaatttga	ttgcacgttt	4980
gtaccaagcg	ggcttatgg	accat	tagagcacac	gcccatatag	acaccaagag	5040
aagagaacca	tacttattca	atgaacctt	gaagtcgata	gtacgtgata	ttatccatt	5100
gagatatttc	ctgctaccta	ccttatacac	catgtttcat	aatcaagtg	tcactggatt	5160
tccgataatg	aatccatgt	ttattgaaca	ccctgaattt	gctgaattgt	atcatatcga	5220
taaccaattt	tactggagta	attcaggct	attagtcaaa	cctgtcacgg	agcctggtca	5280
atcagaaacg	gaaatggttt	tcccacccgg	tatattctat	gaattcgcat	ctttacactc	5340
ttttataaac	aatggactg	atttgataga	aaagaatatt	tctgcacccat	tggataaaat	5400
tccattattt	attgaaggcg	gtcacattat	cactatgaaa	gataagtata	gaagatctt	5460
aatgttaatg	aaaacacgatc	catatgtat	agttatagcc	cctgataccg	agggacgagc	5520
cgttggagat	ctttatgtt	atgatggaga	aactttggc	taccaaagag	gtgagtgacgt	5580
agaaaactcg	ttcattttcg	aaaacaatac	cttaaaaaat	gttcgaatgc	atattccga	5640
gaatttgaca	ggcattcacc	acaatacttt	gaggaatacc	aatattgaaa	aaatcattat	5700
cgcaaaagaat	aatttacaac	acaacataac	gttgaaagac	agtattaaag	tcaaaaaaaaa	5760
tggcgaagaa	agttcattgc	cgactagatc	gtcatatgag	aatgataata	agatcaccat	5820
tcttaaccta	tcgcttgaca	taactgaaga	ttgggaagtt	atttttggc	ccgaacaaaa	5880
actcatctca	gaagaggatc	tgaatagcgc	cgtcgaccat	catcatcatc	atcatttgagt	5940
ttagcctta	gacatgactg	ttcctcagg	caagttggc	acttacgaga	agaccggct	6000
tgctagattc	taatcaagag	gatgtcagaa	tgccatttgc	ctgagagatg	caggcttcat	6060
ttttgatact	tttttatttgc	taacctata	agtataggat	ttttttgtc	attttgtt	6120
ttctcgtacg	agcttgc	tgcgtcc	tgatcagcc	tgcgtat	cttgcgttag	6180
gggttggga	aaatcattcg	agtttgcgt	ttttcttgc	atttccact	cctcttcaga	6240
gtacagaaga	ttaagtgaga	cttcg	tgcggatccc	ccacacacca	tagctcaaa	6300
atgtttctac	tccttttta	cttccaga	tttctcg	ctccgcgc	cgccgtacca	6360
cttcaaaaaca	cccaagcaca	gcataactaa	tttccctct	tttccctct	aggggtcg	6420
taattacccg	tactaaaggt	ttggaaaaga	aaaaagagac	cgcctcg	ttttttctt	6480
gtcgaaaaag	gcaataaaaa	tttttatcac	gtttctttt	cttgaat	tttttttag	6540

-continued

ttttttctc	tttcagtgac	ctccattgtat	attnaagtta	ataaacggtc	ttcaatttct	6600
caagtttcag	tttcattttt	cttggtttat	tacaactttt	tttacttctt	gttcatttaga	6660
aagaaaagcat	agcaatctaa	tctaaggcg	gtgttgacaa	ttaatcatcg	gcatagtata	6720
tcggcatagt	ataatacgcac	aagggtgagga	actaaac			6757

<210> SEQ ID NO 17
<211> LENGTH: 8272
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pAOX2ADE1glslII.

<400> SEQUENCE: 17						
tcgaccggct	gcattaatga	atcgccaaac	gcgcggggag	aggcggtttg	cgtattggc	60
gctttccgc	ttcctcgctc	actgactcgc	tgcgcgttgt	cgttcggctg	cgcgagcgg	120
tatcagctca	ctcaaaggcg	gtataacggt	tatccacaga	atcaggggat	aacgcaggaa	180
agaacatgtg	agcaaaaggc	cagcaaaagg	ccaggaacct	taaaaaggcc	gcgttgctgg	240
cgttttcca	taggctccgc	ccccctgacg	agcatcacaa	aatcgacgc	tcaagtca	300
ggtggcgaaa	cccgacacgga	ctataaagat	accaggcgtt	ccccctgga	agctccctcg	360
tgcgcctctcc	tgttccgacc	ctgccgtta	ccggataacct	gtccgcctt	ctcccttcgg	420
gaagcgtggc	gctttctcat	agtcacgcgt	gtaggtatct	cagttcggtg	taggtcggtc	480
gctccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgcgtc	gccttatccg	540
gtaaactatcg	tcttgagttc	aacccggtaa	gacacgactt	atcgccactg	gcagcagcca	600
ctggtaacag	gattagcaga	gcgaggtatg	taggcgggtc	tacagagttc	ttgaagtgg	660
ggcctaacta	cggctacact	agaaggacag	tatttggtat	ctgcgcctcg	ctgaagccag	720
ttaccttcgg	aaaaagagtt	ggtagcttt	gatccggcaa	acaaaccacc	gctggtagcg	780
gtggttttt	tgttgcag	cagcagatta	cgcgagaaaa	aaaaggatct	caagaagatc	840
ctttgatctt	ttctacgggg	tctgacgtctc	agtggaaacga	aaactcacgt	taagggattt	900
tggcatgag	attatcaaaa	aggatctca	cctagatctt	tttaaatata	aaatgaagtt	960
ttaaatcaat	ctaaagtata	tatgagtaaa	cttggctctga	cagttaccaa	tgcttaatca	1020
gtgaggcacc	tatctcagcg	atctgtctat	ttcggttcatc	catagttgcc	tgactcccc	1080
tcgtgttagat	aactacgata	cgggagggtc	taccatctgg	ccccagtgtc	gcaatgatac	1140
cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	aaaccagcca	gccggaagg	1200
cccgagcgcag	aagtggctt	gcaactttat	ccgcctccat	ccagtctatt	aattgttgcc	1260
gggaagctag	agtaagtagt	tcgcccgtta	atagtttgcg	caacgttgtt	gccattgcta	1320
caggcatcgt	ggtgtcacgc	tcgtcggtt	gtatggcttc	attcagtc	ggttccocaac	1380
gatcaaggcg	agttacatga	tccccatgt	tgtgcaaaaa	agcggtagc	tccttggc	1440
ctccgatcgt	tgtcagaagt	aagttggccg	cagtgttatac	actcatgttt	atggcagcac	1500
tgcataattc	tcttactgtc	atgccatccg	taagatgttt	ttctgtgact	ggtgagttact	1560
caaccaagtc	attctgagaa	tagtgtatgc	ggcgaccgag	ttgctctgc	ccggcgtaa	1620
tacgggataa	tacccgcgeca	catagcgaaaa	ctttaaaagt	gctcatcatt	ggaaaacgtt	1680
cttcggggcg	aaaactctca	aggatcttac	cgctgttgag	atccagtcg	atgtaaaccc	1740
ctcggtgcacc	caactgatct	tcagcatctt	ttacttcac	cagcgttct	gggtgagcaa	1800

-continued

aaacaggaag gcaaaatgcc	gcaaaaaagg gaataaggc	gacacggaaa tggtgaatac	1860
tcataactctt ccttttcaa	tagctccaag gcaacaaatt	gactactcg accgacattc	1920
attcgttatt gattttaat	caacgataaa cggaatggtt	acttgaatga tttcaactta	1980
tgatcattgt ttactaatta	cctaaatagg attttatatg	gaattgaaag aataaggaa	2040
atttcagatg tctgaaaaag	gcgaggaggg tactaatcat	tcaagccat ttcttgcag	2100
taattgcttc ataagcttca	ataactttt cttaactctt	gatagcaatt tctgcattcca	2160
tggctacgcc ctcttgcca	ttaatccgt tggccgtcaa	ccaatctcg agaaaactgct	2220
tatcgtaact ctcttgcgt	ttaccactt ggtaagtctt	ttgattccaa aatctagaag	2280
aatctggagt taaaacttca	tctacttagta ccaattcatt	gttttcgtcc agtccaaatt	2340
cgaatttcgt atcagcaata	atgatcccct taaaaaggc	gaagttttt gcagcagaat	2400
acaactcgac cgcottgaca	gcgaccttct cacaaatgtc	tttacctaca atctcagcag	2460
cttggttcaat agagatgttt	tcatcgtgtt caccctgttc	agctttcggtt gaagggtgtga	2520
aaatcggagt tggaaaggcg	tcgctcttta gaaggttctc	gttttcaacc ttgactccat	2580
ggacagtttt tgagttcttg	tactcttcc atgcacttcc	agtgtatgtt cctctgacaa	2640
tggcttccaa aggtatcagt	ctgtgctttt ttactatcaa	ggatcgccc tctaatttag	2700
atttgtatTT ttcttcagac	agttttgtat gtatgtttagc	aaagacttcc ttgtcattag	2760
aagcaaccaa atgattctt	atgttagggtg caaaaaaatc	aaaccagaaa actgagagct	2820
gagtcaaaat ctccccctta	tcaagatcacatcg	taageggaga	2880
tacggtcagt tgegacgaaac	agcaagttgt tctcatcgac	tgcataaatg tctctaacct	2940
ttccttggc gattaaaggt	aggattccgt ccagatcagt	gttcacaatg gacatacttg	3000
gaaggataca gcaaagtgt	ttgaaagcgta	tgacacatgg aaaggaattt ttcgagttc	3060
cttagatgt atattgggc	ggtgaaagttt	cagatgtta atgcttaata ctcttatact	3120
cttcaaaagcg cccaaatgtt	tctgccaacc	tgactttttt ctgataatg aatcgtaaa	3180
gtggaggatt taaaccatga	ttaagttacg	tgatggcataagg tcgaaaaata	3240
tccgtattca taaacgatta	ttggtaaaag	ttacaaaata ccactaatta cggagaagct	3300
tagtaacagt tatcatctct	tggtcgatttta	acgcttacaa tttccatcg ccattcaggc	3360
tgcgcaactg ttggaaaggcg	cgatcgggtc	gggcctcttc gctattacgc cagggoctcg	3420
aggcacaaac gaacgtctca	cttaatcttc	tgtactctga agaggagtgg gaaataccaa	3480
aaaaaacatc aaactcgaat	gattttccca	aacccctacc acaagatatt catcagctgc	3540
gagataggct gatcaggagc	aagctcgatc	gagaagaaac aaaatgacaa aaaaatcct	3600
atactatata ggttacaat	aaaaaagtat	aaaaatgaa gcctgcattt ctcaggcaaa	3660
tggcattctg acatcctt	gatttataatc	tagcaagacc ggtcttcgt taagtgc	3720
acttgaactg aggaacagtc	atgtctaagg	ctaaaactca atgtatgtt gatgtatgtt	3780
cgacggcgct attcagatcc	tcttctgaga	tgatgtttt ttcggccca aaaataactt	3840
cccaatcttc agttatgtca	agcgataggat	taagaatggt gatcttattt tcatttotcat	3900
atgacgatct agtcggcaat	gaactttctt	cgccattttt tttgacttta atactgtctt	3960
tcaacgttat gttgttgtt	aaattattct	ttgcgataat gatttttca atattggat	4020
tcctcaaagt attgtgggt	atgcctgtca	aattctcgaa aatatgactt cgaacat	4080
ttaaggatatt gtttcgaaa	atgaactgag	tttctacgta ctcacccctt tggtagccaa	4140

-continued

aagtttctcc atcatcaaca taaagatctc caacggctcg tccctcgta tcaggggcta	4200
taactattac atatggatcg ttttcattt acattgaaga tcttctatac ttatcttca	4260
tagtgataat gtgaccgect tcaataaata atggaatttt atccaatggt gcagaaatat	4320
tctttctat caaatcgtt ccattgttt taaaagagtg taaagatgctt aattcataga	4380
atataccggg tggaaaaacc atttccgtt ctgattgacc aggctccgtg acaggttga	4440
ctaatagacc tgaattactc cagtaaaattt ggttatcgat atgataacaat tcagcaaatt	4500
cagggtgttc aataaacattt ggattcatta tcggaaatcc agtgacactt gatttatgaa	4560
acatggtgttta taaggtaggt agcagggaaat atctcaatgt gataatatac cgtactatcg	4620
acttcaaaagg ttcattgaat aagtatggtt ctcttcctt ggtgtctata tggcggtgt	4680
ctctaaaaaa tgggtaccat aagcccgtt ggtaccaacg tgcaatcaat tcaggtgttag	4740
gatcctcagc aaagccagct atgtcggttc ctataatgg cataccagca atgttgc	4800
acagaaccat aggaatggaa atctttaagt aatcccaattt ggccacattt tcaccagtcc	4860
atgtggcagc agtacgttga gagccggcaa aaaaaggccct tgtagaagg aaaggacgct	4920
tatcgatgg tgaataaatac gattttattt cgtcgttaagt agtctcatgc actgatagac	4980
catatatgtt atggacggat ctttcctcaa tggatatttgc tgaaatcaaa tctttggag	5040
ctgtggtctc tggccatcg aaaatcgaag gctcggttcat atcatttcaat atgataaaat	5100
tagttaatac agccggcaga tccataaacc gttcgaaaaa ggacttccaa atctttggc	5160
catatttgct tatggatca atccatatac aattacctgg ccagcaatga cctacatagt	5220
catttcattt gtgatccttgc actgctacat tttcatttacat taccctgtca ctgatttcat	5280
aatctttctt taaatgagga tcgatttagta cgacaagattt tctacccaaat tttttttaatt	5340
tggataacag ctttttggta ttggaaaagg agtgctgtt ccaagttttt tttttttttt	5400
cgttcgatata ctccaagtcc aaccaaataa aatcgtaagg aatcatatga gcatccatct	5460
gagagtccac tggatggatcg tccatctcat cattataattt ccatctacat tgatggatcc	5520
ctatagagga aatgggcggt aaaaagggtt taccagtcaa atcggtttt ttgtcaatga	5580
tagttggaaat atctggcccc agggacatgatc ctacatctat gacaccatttt tcggagatcc	5640
aatgagtcat cgtttttattt ttactgggtt catabtttgc gtctacccaa gtgtcagctg	5700
cattgaccca aaagatagat gtggacgaag atgaaaacat gaatgggatc gaaccgtaca	5760
ttgggtggct ggtaccgtt ttgtactcaa agacatcaac gttgaaaagg ctgttagggtt	5820
cctttccacc tgaagtgtcc atcagccatca ggcacgtcgc atgttccggat ataccgttgc	5880
cattagtaga acccatgaaa gagaaatcta ggcacaccga ttcaggcccc aaaggcatag	5940
agtcatgttt tgaataacaag aaattgtccct taaacatgtt gaaagttttt tcttctggca	6000
gcacgtgtgc gaagtttcc tgcgttgc tatgtatgtt aatgttccagg aaattttgtt	6060
cgtttacaat aagtttccgc gcattttgcg agttaactttt caattggaaaa ggttcggcaaa	6120
agatttctac ggatacatca ccgtttcgaa gatgaaatgt gtctcagtg gagtttggaaa	6180
gtgacaaaaaa tgaagatattt ttgcaccaga atgagttcac agtttggttt tgcttaagga	6240
agtggaaattt tggatatactg gtcctgttcg ctttccttgc aaatttcttgc tgcgttgc	6300
acttccaggt ctcattgttgc acgttgcgttgc agatcaacaa accgctgttgc ttgggtggca	6360
ttcttcctttt ctcattttata gtgaaaccttgc ctgactgttgc ctgttttttggaaaaa	6420
atggaaactg aacggctata tcatcgccctt ccaatcttgc tatagttttta attatgttag	6480
catgaagcac attctctaaa ggatcgtgttgc caatagactc ggcgtccact ttgttaatgc	6540

US 9,359,628 B2

49

50

-continued

agtgtatgaga tttggacaata tttctgcataaaccctgtt tctatggcaaaacccagatt
gcccacactt cttaataga tagtcgttaa acgcatacgaaaaggcgta aagaagacca
atggcatac gagccatttc aaaaggacca tctcgaggta ccgatccgag acggccggct
ggccacgtg aattttttc tcagttgatt tgtttggtgg gathtagtaa gtctaaact
tcgttaaaaa agatcaatgt agtcaatacagttgatccga aatagaagga agaggttgc
aatgtgtaaag aacaatgttag taaaagccc gtttaaagac aatattctt gatgtgtatc
agaaaaggac aataaggat ttgggtgtct tcttttatac caataatcgctccatcatcg
cttaatttc tccccatctc aaccgggtaa gggttaggacg cttctgtataatgttcacat
aaaagggtt ttcactccga gacaaaatt tatgcgacaa aaatagctatcttggaaagg
tcatgttta tcaacttgca ttgttgcaaa ggagaagcaa ggacaactca acatggtaa
aaattcaaaa ccaaccaatt gggaaactccc aactgtccac tagtagtgc acagctgtca
cttttgcgt tctgtgtctt gtcttttgccttaactgtc ctcttacggctttttcc
aaaaaatcat tggggaaatgt tgccctcatcagagtccaa tgaccatga ataaagttc
ttgtactgtt taagacgtat aattgcaacg ataatccgag cagttacgg ggtacatcac
gtgcttgca tatgatctcg gagtcggatc agttccggat gtgtatgtt accccatagt
ttcaactct aatgcggcc ccaagtggca tacaccctcc atcaatcttatcttggaaagg
tttaccatc gttgggtggat gatgatgact cgcttagtct ctgctgttcg atattaactt
tgtaaggatc gcccggat gggaaattga ggggttgtaa cctgaatttg caggctactt
acattggact tttgagaagg ctggacgggt gatgaagagg gctgggtgca gggaaatgg
aaaaaatcta gttgagagga ctgttggaaa tttagggaaa tggagtccttaaactgtgaca
aaactcaag gatggggatt ttcatgtac ttgttcatgc ctgcacaaag ctaaaggaag
gtatttgatt ctggataaaat ggatattgtatctgtttag cagatgtca agttctacta
gtatgtct ggtatctcgat agccttcaat tggcgatc ttactcgaaatgttattt
ttatgtac agacgaagaa cgagagatgttgcacatt cagaggtaaag acaatatgtc
gtattatcaa aataagtatac gaaccttatc tagggatcctatcgatccaaatgtca
tagtgggtat tgcgttgcataatctgtatcgt gtttggagatgttgcgtat
gatgttttagt aaatgtttt cagaaaaggc gctaaatggaaa taaccatgtaa gtaataaata
acgtaaaatgttgcataatcaaaagaat caagatagag gtcaaaagcat agataatccc
cccgtagaaatggaccggcatatgttgcataaaaaaaaatgttgcataatgtca
8272

```
<210> SEQ ID NO 18
<211> LENGTH: 5727
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of
      plasmid pGAPZAGLSSII.
```

<400> SEQUENCE: 18

tcggagatggt ccttttggaaa ttggctcgat gccaatttgtt cttctttacc gctttttcgc	60
atgcgtttac cgactatcta tttaagaagt gtgcgcaatc tgggtttgc catagaaaca	120
gggtttatgc agaaaatatt gccaaatctc atcactgtca ttacaaagtg gacgcgcgat	180
ctattgcaca cgatcctta gagaatgtgc ttcatgtcac cataataaa actataccaa	240
gattggaggcgatgatata gcccgttcgt tcccatgttc tctctttttt ttacaggatc	300

-continued

actcagtaag gttcaactaa aatgagaaaa agagaatgcc aaccaacagc agcggttgt 360
tgatctttc acaacggtc aatgagacct ggaagtacgc attcgacaag aaatttcaag 420
aggaggcga caggaccagt attccacaat tccacttcc taagaaaaaa caaactgtga 480
actcattctg gtcgaaaata tcttcattt tgtcaacttc aaactccact gcagacacat 540
ttcatctcg aaacggtgat gtatccgtag aaatcttgc tgaaccttt caattgaaag 600
tttactggca aatcgctg aaacttattt taaacgagca aaatttctg aacatgAAC 660
atcatagaac taagcaggaa aacttcgcac acgtgctgcc agaagaaaaca actttcaaca 720
tgtttaagga caatttctg tattcaaagc atgactctat gccttgggg cctgaatcg 780
tttgcgtaga tttcttttc atgggttcta ctaatgtcta cggtataccg gaatcgca 840
cgtcgctaag gctgtggac acttcaggta gaaaggAAC ctacaggctt ttcaacgttg 900
atgtcttgc gtacaacatc ggtaccagcc aaccaatgtc cggttcgtat ccattcatgt 960
tttcatcttc gtccacatct atctttggg tcaatgcAGC tgacacttgg gtagacataa 1020
agtatgacac cagtaaaaat aaaacgtga ctcattggat ctccgaaaat ggtgtcatag 1080
atgtagtcat gtccctgggg ccagatattc caactatcat tgacaattt accgatttga 1140
ctggtagacc ctttttaccc cccatttctt ctataggta ccatcaatgt agatggaatt 1200
ataatgtga gatggacgtt ctcacagtgg actctcagat ggatgtcat atgatccct 1260
acgattttat ttgggtggac ttggagtata cgaacgacaa aaaatattt acttggaaAGC 1320
agcactctt tccccatcca aaaaggctgt tatccaaatt aaaaaagttt ggtagaatAC 1380
ttgtcgtaat aatcgatctt cattaaaga aagatttgc aatcgtgtc agggtaatta 1440
atgaaaatgt agcagtcaag gatcacaatg gaaatgacta tggatgtcat tgctggccag 1500
gtatttat atggattgtat accataagca aatatggccaa aagatttgg aagtccccc 1560
tcgaacgggt tatggatctg cgggtgatt taactaattt attcatttgg aatgatgt 1620
acgagccttc gatTTcgat ggcccagaga ccacagctcc aaaagatttgg attcacgaca 1680
attacattga ggaaagatcc gtcataaca tatatggct atcgtgtcat gaagctactt 1740
acgacgcaat aaaatcgatt tattcaccat ccgataagcg tcctttccctt ctaacaagg 1800
cttttttgc cggctctcaa cgtactgtg ccacatggac tgggtgacaaat gtggccaaatt 1860
gggattactt aaagattttcc attcttatgg ttctgtcaaa caacattgtt ggtatggcat 1920
ttataggagc cgacatagct ggcttgcgtg aggatccatc acctgttgcattt attgcacgtt 1980
ggtaccaagc gggcttatgg tacccattttt ttagagcaca cgcccatata gacaccaaga 2040
gaagagaacc atacttattt aatgaaccc tgaagtgcgt agtacgtgtat attatccaaat 2100
tgagatattt cctgtaccc accttataca ccatgtttca taaatcaagt gtcactggat 2160
ttccgataat gaatccaaat tttattgtaac accctgttgcattt tgctgttgcattt tatcatatcg 2220
ataaccaatt ttactggagt aattcagggtc tattagtcaat acctgtcactt gagectggtc 2280
aatcagaaac ggaaatgggtt ttcccaccccg gtatattctt tgaattcgca tctttacact 2340
cttttataaa caatggtaact gatttgcattt aaaaagatattt ttctgttgcacca ttggataaaaa 2400
ttccattttt tattgttgc ggtcacatca tcactatgaa agataagtat agaagatctt 2460
caatgttaat gaaaaacgtat ccatgttgcattt tagttatgc ccctgttgcattt gagggacgag 2520
ccgttggaga tctttatgtt gatgtggag aacttttgg ctaccaaaaga ggtgagttacg 2580
tagaaactca gttcattttc gaaaacaata ccttaaaaaaa tggatgttgcattt catattcccg 2640

-continued

agaatttgac	aggcattcac	cacaatactt	tgaggaatac	caatattgaa	aaaatcatta	2700
tcgcaaagaa	taatttacaa	cacaacataa	cgttgaaaga	cagtattaaa	gtcaaaaaaa	2760
atggcgaaga	aagttcattt	ccgactagat	cgtcatatga	aatgataat	aagatcacca	2820
ttcttaacct	atcgcttgc	ataactgaag	attggaaat	tatttttggg	cccgaaacaaa	2880
aactcatctc	agaagaggat	ctgaatagcg	ccgtcgacca	tcatcatcat	catcatggag	2940
tttttagcctt	agacatgact	gttcctcagt	tcaagttggg	cacttacgag	aagaccggtc	3000
ttgcttagatt	ctaatacaga	ggatgtcaga	atgccatttgc	cctgagagat	gcaggottca	3060
tttttgatac	tttttttattt	gtaacctata	tagtatagga	tttttttgtt	cattttgttt	3120
cttctcgatc	gagcttgc	ctgatcagcc	tatctcgac	ctgatgata	tcttggttgc	3180
ggggtttggg	aaaatcattt	gagtttgc	tttttcttgg	tatcccac	tcctcttcag	3240
agtacagaag	attaagtgc	accttcgtt	gtgcggatcc	cccacacacc	atagttcaa	3300
aatgtttcta	ctccctttt	actcttccag	attttctcgg	actccgcgc	tcgcgcgtacc	3360
acttcaaaac	acccaagcac	agcatactaa	atttcccttc	tttcttcgtt	taggggtc	3420
ttaatttaccc	gtactaaagg	tttgaaaag	aaaaaaagaga	ccgcctcg	tctttttttt	3480
cgtcgaaaaaa	ggcaataaaaa	attttatca	cgtttctttt	tcttgaattt	ttttttttta	3540
tttttttctt	ctttcagtga	cctccattga	tatthaagtt	aataaacgg	cttcaatttc	3600
tcaagtttca	gtttcatttt	tcttgcgttca	ttacaacttt	ttttacttct	tgttcttag	3660
aaagaaagca	tagcaatcta	atctaagggc	ggtggttgc	attaatcatc	ggcatagttat	3720
atcggcatag	tataatacga	caaggttgg	aactaaacca	tggccaagtt	gaccagtgc	3780
gttccgggtgc	tcaccgcgc	cgacgtcgcc	ggagcggcgt	agttctggac	cgaccggc	3840
gggttctccc	gggacttcgt	ggaggacgac	ttcgcgggt	tggtccggg	cgacgtgacc	3900
ctgttcatca	gcgcgggtca	ggaccaggt	gtgcggaca	acaccctggc	ctgggtgtgg	3960
gtgcgcggcc	tggacgagct	gtacgcccag	tggtgcggagg	tcgtgtccac	gaacttccgg	4020
gacgcctccg	ggccggccat	gaccgagatc	ggcgagcagc	cgtggggcgc	ggagttcgcc	4080
ctgcgcgacc	cgccggccaa	ctgcgtgcac	ttcgtggccg	aggagcagga	ctgacacgtc	4140
cgacggccgc	ccacgggtcc	caggcctcg	agatccgtcc	cccttttcc	ttgtcgatat	4200
catgttattt	gttatgtcac	gttttacattt	acgcctcccc	cccacatccg	ctctaaccga	4260
aaaggaagga	gttagacaac	ctgaagtcta	ggtccctattt	tatTTTTTta	tagttatgtt	4320
agtattttaa	acgttattttt	tatTTcaat	tttttctttt	tttctgtaca	gacgcgtgt	4380
cgcatgtaac	attatactga	aaaccttgc	tgagaagg	ttggggacgt	cgaaggctt	4440
aatttgc	ctggagacca	acatgttgc	aaaaggccag	caaaggccca	ggaaccgtaa	4500
aaaggccgc	ttgtctggcgt	ttttccatag	gtccgc	cctgacgac	atcacaaaaaa	4560
tcgcgtca	agtcagaggt	ggcgaaaccc	gacaggacta	taaagatacc	aggcggttcc	4620
cccttggaa	tccctcg	gtctctctgt	tccgaccc	ccgcttac	gataccgtc	4680
cgccttctc	ccttcggaa	gctgtggcgt	ttctcaatgc	tcacgtgt	ggtatctcag	4740
ttcggtgt	gtcggtcg	ccaagctgg	ctgtgtgcac	gaaccccc	tgcagccg	4800
ccgctgcgc	ttatccggta	actatcg	tgagtccaa	ccggtaagac	acgacttac	4860
gccactggca	cgagccactg	gtacacaggat	tagcagagcg	aggtatgt	gcgggtgtac	4920
agagttctt	aagtgggtgc	ctaaactacgg	ctacactaga	aggacagtt	ttggtatct	4980
cgctctgt	aagccagtt	ccttcggaaa	aagagttgg	agcttgc	ccggcaaaaca	5040

-continued

```

aaccaccgct ggtagcggtg gttttttgt ttgcaagcag cagattacgc gcagaaaaaa 5100
aggatctcaa gaagatcctt tgcgttttc tacggggct gacgctca ggaacgaaaa 5160
ctcacgttaa gggatttgg tcatgcata gatcagatct ttttgtaga aatgttttgg 5220
tgtcctcgta caatcaggtt gccatctctg aaatatctgg ctccgttgc actccgaacg 5280
acctgtggc aacgtaaaat tctccgggtt aaaacttaaa tgtggagtaa tggaaccaga 5340
aacgtctttt cccttcttc tccttccacc gcccgttacc gtccctagga aattttactc 5400
tgctggagag ctctttctac ggcccccttg cagcaatgtt ctccctagca ttacgttgc 5460
gtaaaaacgg aggtcggtt cccgacccat cagcccaggat atggaaaagt cccggccgtc 5520
gctggcaata atagcggggc gacgcattgtc atgagattat tgaaaccac cagaatcgaa 5580
tataaaaggc gaacacccctt cccaattttt gtttctctg acccaaagac tttaaattta 5640
atttatgtt ccctatttca atcaattgaa caactatttca gaaacgagga attcaatgtt 5700
cccagccggc cgtctcgat cggtacc 5727

```

<210> SEQ ID NO 19
<211> LENGTH: 7236
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pGAPADE1gl1sII.

<400> SEQUENCE: 19

```

tcgaccggct gcattaatga atcggccaac gcgccccggag aggccgttttgcgtattggc 60
gctttccgc ttcttcgttc actgacttcgc tgccgtcggt cggtccgtcg cggcgagcgg 120
tatcagctca ctcaaaggcg gtaataacggt tatccacaga atcaggggat aacgcaggaa 180
agaacatgtg agcaaaaggc cagcaaaagg ccaggaaccc taaaaaggcc gcgttgcgtt 240
cgttttcca taggctccgc cccctgcacg agcatcacaa aaatcgacgc tcaagtccaga 300
ggtggcgaaa cccgacagga ctataaagat accaggcggtt tccccctggaa agctccctcg 360
tgcgtctcc tggccgttcc ctggatcacct gtccgcctttt ctcccttcgg 420
gaagcgtggc gctttctat agtcacgcgt gtaggtatct cagttcggtt taggtcggtt 480
gctccaagct gggctgtgtg cacgaaccccc ccgttcagcc cgaccgcgtc gccttatccg 540
gtaactatcg tcttgagttcc aacccggtaa gacacgactt atcgccactg gcagcagcc 600
ctggtaacag gattagcaga gcgaggatgtt taggcgggtgc tacagagttc ttgaagtgg 660
ggcctaacta cggctacact agaaggacag tattttgtat ctgcgtctcg ctgaagccag 720
ttaccttcgg aaaaagagtt ggtagctttt gatccggcaaa acaaaccacc gctggtagcg 780
gtggttttt tggccgttcc cggcgatccaa cggcgacggaaa aaaaggatctt caagaagatc 840
ctttgtatctt ttctacgggg tctgacgttc agtggaaacgcg aaactcacgt taagggtttt 900
tggtcatgg attatcaaa aggtatctca cctagatctt tttaaatcaa aaatgaagtt 960
ttaaatcaat ctaaaagtata tatggataaa ctgggtctga cagttacca tgcttaatca 1020
gtgaggcacc tatctcagcg atctgtctat ttccgttcatc catagttgcc tgactcccc 1080
tcgtgttagat aactacgata cggggggct taccatctgg ccccaatgtt gcaatgatac 1140
cgccgaccc acgttcacccg gtcggatattt tttttttttt aaaccaggccca gccggaaagg 1200
ccgagcgcag aagtgggtt gcaactttat ccgcctccat ccgttattt aattttttttt 1260
ggaaagctag agtaatgtt tggccgttta atagtttgcg caacgttgcg gccattgtca 1320

```

-continued

caggcatcgt ggtgtcacgc tcgtcgttt gatatggcttc attcagctcc ggttcccaac	1380
gatcaaggcg agttacatga tccccatgt tggcaaaaa agcgggttagc tccttcggtc	1440
ctccgatcgt tgctagaagt aagttggccg cagtgttatac actcatggtt atggcagcac	1500
tgcataattc tcttactgtc atgccatccg taagatgctt ttctgtgact ggtgagtaact	1560
caaccaagtc attctgagaa tagtgtatgc ggcgaccgag ttgcttgc ccggcgtaa	1620
tacgggataa taccgcgcca catagcgaaa cttaaaaagt gctcatcatt ggaaaacgtt	1680
cttcggggcg aaaactctca aggatcttac cgctgtttag atccagttcg atgtAACCC	1740
ctcgtgcacc caactgatct tcagcatctt ttactttcac cagcgttct gggtgagcaa	1800
aaacaggaag gcaaaatgcc gcaaaaaagg gaataaggc gacacggaaa tggtgaatac	1860
tcataactctt ccttttcaa tagctccaag gcaacaaatt gactactcg accgacattc	1920
attcgttatt gatTTAAAT caacgataaa cggaatggtt acttgaatga ttcaCTTAA	1980
tgatcattgt ttactaatta cctaaatagg attttatatg gaattggaa aataaggaa	2040
atTCAGATG tctgaaaaag gcggggggg tactaatcat tcaagccat ttcttgcag	2100
taattgcttc ataagcttca atatactttt cttaacttctt gatacgaaatt tctgcattca	2160
tggctacgcc ctcttgcaca ttcaatccgt tggccgtcaa ccaatctcg agaaaactgct	2220
tatcgtaaact ctcttgcgtat ttacccactt ggtaagtctt ttgattccaa aatctagaag	2280
aatctggagt taaaacttca tctacttagta ccaattcatt gtttctgtcc agtccaaatt	2340
cgaatttcgt atcagcaata atgatccc tcaaaaggc gaagttttt gcagecagaat	2400
acaactcgac cgccttgaca gegacccctt cacaatgtc ttacccatca atctcagcag	2460
cttgcattcaat agagatgttt tcatcggtt caccctgttc agcttcgtt gaaggtgtga	2520
aaatcggagt tggaaaggcg tegctcttctt gaagggtctc gtttcaacc ttgactccat	2580
ggacagtttt tgagttcttg tactcttcc atgcacttcc agtgtatgtt cctctgacaa	2640
tggcttccaa aggtatcgt ctgtgccttt ttactatcaa ggatcgccc tctaatttag	2700
atTTGTATTt ttcttcagac agttttatgtt gtagtaaagc aaagacttcc ttgtcattag	2760
aagcaaccaa atgattctt atgttaggggtt ccaaaaaatc aaaccagaaa actgagagct	2820
gagtcaaaat ctcccctta tcaagaaatac cgtttgcatt aatcacatcg taageggaga	2880
tacggtcagt tgccgacgaa acgaaatgtt tctcatcgac tgcataatcg tctctaacct	2940
ttccttggc gattaaaggt aggatccgtt ccagatcgtt gttcacaatg gacatacttg	3000
gaaggataca gcaaaatgtt tggaaaggcg tgacacatgg aaaggaattt ttggatTTT	3060
ctagatgtt atattggggc ggtgaaaggtt cagatgttta atgcttaata ctcttatact	3120
cttcaaagcg cccaaatgtt tctgccaacc tgactttttt ctgataatcg aatcgtaaa	3180
gtggatgtt taaaccatga ttaagttacg tgatTTGGCA ctggataagg tcgaaaaata	3240
tccgtattca taaacgatttta ttggtaaaag ttacaaaata ccaactaatta cggagaagct	3300
tagtaacagt tatcatctt tggtcgatca acgcttacaa ttccattcg ccattcaggc	3360
tgcgcacactg ttggaaaggcg cgatcggtgc gggcccttgc gctattacgc caggccctcg	3420
aggcacaaac gaaacgtctca cttaatcttc tgtaactctga agaggatgg gaaataccaa	3480
aaaaacatc aaactcgaat gatTTCCCA aacccttacc acaagatatt catcagctgc	3540
gagataggct gatcaggagc aagctcgatc gagaagaaac aaaatgacaa aaaaatcct	3600
atactatata ggttacaaat aaaaaagtat caaaaatgaa gcctgcattt ctcaggccaa	3660

-continued

tggcattctg acatccttta gattagaatc tagcaagacc ggtttctcg taagtgcaca	3720
acttgaactg aggaacagtc atgtctaagg ctaaaactca atgatgtatgatgtatgg	3780
cgaaggcgctt attcagatcc tcttctgaga tgagtttttgc ttcggggccca aaaataactt	3840
ccaaatcttc agttatgtca agcgataggta aagaatggt gatcttattatcatttcatt	3900
atgacgatct agtcggcaat gaactttctt cgccattttt tttgacttta atactgtctt	3960
tcaacgttat gttgtgttgtt aaattattct ttgcgataat gatttttca atattggat	4020
tcctcaaagt attgtggta atgcctgtca aattctcgaa aatatgactt cgaacatttt	4080
ttaaggattt gtttcgaaa atgaactgag tttctacgtt ctcacccctt tggtagccaa	4140
aagtttctcc atcatcaaca taaagatctc caacggctcg tccctcggtt tcaggggctt	4200
taactattac atatggatcg ttttcttta acattgaaga tcttctataat ttatcttca	4260
tagtgataat gtgaccgcct tcaataaataa atgaaattttt atccaaatggt gcagaaatata	4320
tctttcttat caaatcagta ccattgttta taaaagatggt taaagatgcg aattcataga	4380
atataccggg tgggaaaacc atttccgttt ctgattgacc aggctccgtg acagggttga	4440
ctaataagacc tgaattactc cagtaaaattt ggttatcgat atgatataat tcagcaaattt	4500
cagggtgttc aataaacattt ggattcattt tccggaaatcc agtgcacattt gatattgaa	4560
acatgggtgtttaaaggtaggtt agcaggaaat atctcaattt gataatataatcgcgttactatcg	4620
acttcaaaagg ttcattgtttaaattt aagtatggttt ctcttctttt ggtgtctata tggcggtgtt	4680
ctctaaaaaa tgggttaccat aagcccgtt ggtaccaacg tgcaatcaat tcaggtgttag	4740
gatcctcagc aaagccagct atgtcggctc ctataatgg cataccagca atgtgttttgc	4800
acagaaccat aggaatggaa atctttaatgtt aatccaaattt ggccacatttgc accaggatcc	4860
atgtggcagc agtacgtttaaaggccctt gttttagaagg aaaggacgct	4920
tatcggtatgg tgaataaattt gattttatttgc gtcgttaatgc actgatagac	4980
catatatgtt atggacggat cttcccttcaat tgtaatttgcgtt gtaatcaaaat tcttttggat	5040
ctgtggcttc tggccatcg aaaatcgaag gtcgttcat atcatttcaat atgatataat	5100
tagttaatc agccggcaga tccatataacc gttcgaaaaaa ggacttccaaatcttttggc	5160
catatttgc tatggatca atccatataatgaaatttgcgtt ccagcaatgatcctacatagt	5220
cattttccattt gtgatcccttgc actgttcatat ttccatataat taccctgtca ctgttcat	5280
aatctttttttaatgatggatc tcgatttagta cgacaagattt tcttccaaat tttttttaatt	5340
tggataacag cttttttggat tgggaaaagg agtgctgtt ccaagtttttgc	5400
cgttcgtata ctccaagtcc aaccaatataa aatcgtaagg aatcatatgatc gcatccatct	5460
gagagtccac tggatggatc tccatctcat cattataattt ccatctcatat tgatggatcc	5520
ctatagagga aatggggcggtt aaaaagggtt taccgttcaat atcgtaatgc ttgtcaatgc	5580
tagttggat atctggcccccc agggacatgatc ctacatctat gacaccattt tgggatcc	5640
aatgagtcat cgttttatttttactggatc catactttat gtctaccatgc gtgtcagct	5700
cattgaccca aaagatagat gtggacgaaatgatc atgaaaacat gatggatc gaaccgtaca	5760
ttgggtggctt ggtaccgttgc ttgtacttcaat agacatcaac gttgaaaaggc ctgttagggat	5820
cctttccacc tgaagtgtcc atcagccttgc ggcacgttcgc atgttccgtt ataccgttgc	5880
cattagtaga acccatgaaa gagaaatgttcaat ggcacaccgc ttcaggcccccaaaaggatcg	5940
agtcatgtt tgaatacaatgaaatttgcctt taaacatgtt gaaatggat tcttctggca	6000
gcacgtgtgc gaagtttcc tgcgttgc tatgtatgttc aatgttcagg aaattttgc	6060

-continued

cgtttacaat aagtttcagc gcattttgcc agtaaaactt caattgaaaa ggttcagcaa	6120
agatttctac ggatacatca ccgtttcgaa gatgaaatgt gtctgcagt gагtttggaaa	6180
gtgacaaaaaa tgaagatatt ttgcaccaga atgagttcac agtttggttt tgcttaagga	6240
agtggaaattt tggaataactg gtcctgttcg ctcctcttg aaatttcttg tcgaatgcgt	6300
acttccaggt ctcattgaac cggtgtgaag agatcaacaa accgctgctg ttgggtggca	6360
ttctctctt ctcatatata gtgaacctta ctgagtgtc ctgtaaaaaa gagagagaga	6420
atgggaactg aacggctata tcatgcctt ccaatcttgg tatagttta attatggtag	6480
catgaagcac attctctaaa ggatcgtgt caatagactc ggcgtccact ttgtataagc	6540
agtgtatgaga tttggcaata tttctgtcat aaaccctgtt tctatggcaa aaccaggatt	6600
ggcacacactt cttaataga tagtcggtaa acgcgcgcga aaaagcggtt aagaagacca	6660
attggcatac gagccatttc aaaaggacca ttcgcgggtt ccgcgcgcgc acggccggct	6720
ggccacacgtg aattccctgtt ttgcggaaatag ttgttcaatt gattgaaata gggacaata	6780
aattaaattt aaagtctttt gggtcaggaga aacccaaatt gggaaagggtt ttgcctttt	6840
atattcgatt ctgggtggttt ccaataatct catgacatgc gtccgcgcgc tattattgcc	6900
agcgacggcc gggacttttc catccctggg ctgcgtggc gggtacacga ctcgggtttt	6960
acccgcacacg taatgcgtggg aagagcattt ctgcgggggg gcccgtttag aagctctcca	7020
gcagagtaaa atttccttagg gacggtaacg ggccgtggaa ggagagagaa gggaaagagac	7080
gtttctgggtt ccattactcc acatattaatgt ttaccccg agaattttac gttgcagca	7140
ggtcgttcgg agttgcaacg gagccagata ttccagatgg ggcgttgcgaa ttggacagg	7200
acaccaagac atttctacaa aaaagatctg atctca	7236

<210> SEQ ID NO 20
 <211> LENGTH: 6173
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pPICZAGL5II.

<400> SEQUENCE: 20

cgaacaaaaaa ctcatctcg aagaggatct gaatagcgcc gtcgaccatc atcatcatca	60
tcattgagtt tggtagcctta gacatgactg ttccctcgat caagttgggc acttacgaga	120
agaccgggtct tgcgttagattc taatcaagag gatgtcagaa tgccatttgc ctgagatgt	180
caggcttcat ttttgataact tttttatggtaaacctat agtataaggat tttttttgtc	240
attttggtttc ttctcgatcg agtttgcgtcc tgatcagcct atctcgacgc tgatgaatat	300
cttgcgttag gggtttggga aaatcattcg agtttgcgtt ttttcttggat atttccacact	360
cctcttcaga gtacagaaga ttaagtggaa ctttcgtttg tgccgttccccc ccacacacca	420
tagttcaaa atgtttctac tccttttttca ctcttcaga ttttctcgga ctccgcgcgt	480
cggccgttacca cttcaaaaaca cccaaagcaca gcatactaaa ttttccctct ttcttcctct	540
agggtgtcgtaatttcccg tactaaagggt ttggaaaaga aaaaagagac cgcctcgat	600
ctttttcttc gtcgaaaaag gcaataaaaaa tttttatcac gtttctttt cttgaaattt	660
ttttttttag ttttttttc ttccgtgc ctccattgtat atttaagttt ataaacggtc	720
ttcaatttctt caagtttcag tttcatat tttgttctat tacaactttt ttacttctt	780
gttcattttaga aagaaagcat agcaatctaa tctaaggggc ggtgttgaca attaatcatc	840

-continued

-continued

ttccaaattgttcaagcttttgcattttaaacgacttttaacgacaacttgagaagatcaaaaa 3240
acaactaattatcgaaacaggaaatttcacgtggccccagccggcgctcggatcggtac 3300
ctcgagatggtcctttgaaatggctcgtagtgccaaattggcttttaccgctttcg 3360
catgcgttaccggactatctattnaagaagtgtgcgcaatctggggtttccatagaaac 3420
agggtttatcagaaaatattgccaatctcatcactgctattacaaagtggacgcccag 3480
tctatttgcacacgttcctttagagaatgtgttccatgtctccataattaaactatacca 3540
agattggaggcgcatgtatagccgtttccattctctctttttacaggat 3600
cactcgttaagggttcaacttaaatgagaaaagagaaatgccaaccaacagcagcggttt 3660
ttgatcttccacaacggttcaatgagacc tggaagtacgcattcgacaa gaaatttcaa 3720
gaggaggcgaacaggaccagtatccacaa ttccacttcc ttaagcaaaaacaactgtg 3780
aactcattctggtcgaaaatatcttcattttgtcactttcaaactccac tgcagacaca 3840
tttcatcttgcggatcgatgtatccgtatcc ttaagcaaaaacaactgtg 3900
gtttactggcaaaatgcgtgaaacttattgtaaacgagc aaaatttctgaacattgaa 3960
catcatagaactaaggcggaaacttcgcacgtgtgc cagaagaaac aactttcaac 4020
atgttaaggacaatttcttgtattcaagcatgactcta tgcctttgggcctgaatcg 4080
gttgcgttagatttcttccatgggttctactaatgttacggatataccgaaacatcg 4140
acgtcgctaa ggctgtatggacacttcaggtggaaaggAACccacaggctttcaacgtt 4200
gatgtctttagtacaacatcggttccatgggttctactaatgttacggatataccgaaacatcg 4260
tttcatcttcgtccacatctatctttgggtcaatgtcactgttgcacacttggtagacata 4320
aagtatgaca ccagtaaaaaataaaacgtatctcattggatctccggaaaa tggtagtcata 4380
gatgtatca tggccctggggccagatattccaactatcta ttgacaaatttaccgattt 4440
actggtagaccccttttaccgccttccatgggttctatagggtaccataatgttagatggat 4500
tataatgtatctatggacgttctcacagtgactctcaga tggatgtctatgtatttctt 4560
tacgattttatgttgggttgccttgggtatctacggacacaaaaaatattttacttggaa 4620
cagcacttcttccatccaaaaaggctgttatccaaataaaaaagttggtagaaat 4680
cttgcgttacatcgatccatctttaagaaagattatgaaatcgatgtacgggttac 4740
aatgaaaatgtacgttcaaggatcacaatggaaatgtactatgttca tggctggca 4800
ggtaattctatggattgtatccatatacgaaatatgtggccaaatgttgcctt 4860
ttcgaacggttatggatctgggttctatggatcttactaatttatttgcgttac 4920
aacgagcccttcgtatccatgggttctatggatcttactaatttatttgcgttac 4980
aattacatttggggaaagatctgtccataacatataatgttacgtgtacgttac 5040
tacgacgcataaaaaatgtatccatgggttctatggatcttactaatttatttgcgttac 5100
gcttttttgcgttacatggatcttactaatttatttgcgttac 5160
tgggattacttaaagatttcatccatgttgcgttacatggatcttactaatttatttgcgttac 5220
tttatagtaggcggacatagtcggcttgcgttacttgcgttacatggatcttactaatttatttgcgttac 5280
tggtagccatggggctttagtgcgttacttgcgttacatggatcttactaatttatttgcgttac 5340
agaagagaacatacttattcaatgtacgttacttgcgttacatggatcttactaatttatttgcgttac 5400
ttgagatatttcctgttaccatccatgttacttgcgttacatggatcttactaatttatttgcgttac 5460
tttccgataatgtacgttacttgcgttacatggatcttactaatttatttgcgttac 5520
gataaccaatttactggatcttgcgttacatggatcttactaatttatttgcgttac 5580

-continued

```

caatcagaaa cgaaaaatggt tttcccaccc ggtatattct atgaattcgc atcttacac      5640
tctttataa acaatggta tgatttgata gaaaagaata tttctgcacc attggataaa      5700
attccattat ttattgaagg cggtcacatt atcactatga aagataagta tagaagatct      5760
tcaatgttaa tggaaaacga tccatatgtat agatttatag cccctgatac cgagggacga      5820
gccgttggag atctttatgt tgatgtatggaa gaaacttttgcgtacccaaag aggtgagttac      5880
gtagaaactc agttcatttt cggaaaacaat accttaaaaaa atgttgcagatcatatccc      5940
gagaatttga caggcattca ccacaataact ttgaggaata ccaatattga aaaaatcatt      6000
atcgcaaaaga ataatttaca acacaacata acgttgcagatcatatggataa agtcaaaaaaa      6060
aatggcgaag aaagttcatt gcccactaga tcgtcatatg agaatgataa taagatcacc      6120
attcttaacc tatcgcttga cataactgaa gattggaaag ttatTTTGG gcc                  6173

```

```

<210> SEQ ID NO 21
<211> LENGTH: 7639
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of
plasmid pPICADE1glsII.

```

```

<400> SEQUENCE: 21
aaattccctcg tttcgaataa ttagttgttt tttgatcttc tcaagttgtc gttaaaatgc      60
gttaaaatca aaagcttgc aatttggaaacc agtcgcattt atgaaaatgtt gctaataatg      120
atgataaaaaa aaaagggtta agacaggcgc gcttccttct gtttatatat tgctgtcaag      180
taggggttag aacagttaaa ttttgcattt gaacgtttagg ctatcagcag tattccacc      240
agaatcttgg aagcatacaa tggcggagaca atgcataatc atccaaaaag cgggtgtttc      300
cccatttgcg tttcggcaca ggtgcaccgg gggtcagaag cgatagagag actgcgcctaa      360
gcattaatga gattatttt gggcattcgtt caatcaatac caaacaagac aaacggatgt      420
ccgacttttg gaagtttctt tttgaccaac tggccgttag catttcaacg aaccaaactt      480
agttcatctt ggttgcgcgc acgtttttgtt catatttagt tccaagacag cgttttaactt      540
gtcagttttt ggccatttgg ggaacatgaa actatggatcccacactca gaaaggccctc      600
atctggatgtt atgttgcgggtt gtaatgcggat gtttgcgcgc ttcggaaata aacaaacatg      660
aacctcgcca gggggggccag gatagacagg ctaataaaatgtt catgggttta gtagcctaat      720
agaagggattt ggaatgagcg agtccaaatc aagcccaata actgggttgcg tttttcgatg      780
gcaaaaatgtt gtttgcgcgc gaaagggatgtt gggttgcgcgc ggttgcgcgc ggttgcgcgc      840
tagtgtatcc cttttttttttt cttttttttttt cttttttttttt cttttttttttt cttttttttttt      900
gatggcaaaa aggtttcattt caacctttcg tttttttttttt tttttttttttt tttttttttttt      960
tgaatcgccca aacggccgggg gagagggcgat tttttttttttt tttttttttttt tttttttttttt      1020
ctcaactgact cttttttttttt cttttttttttt cttttttttttt cttttttttttt cttttttttttt      1080
gcggtaatac ggttatccac agaatcaggat gataacgcag gaaagaacat gtgagcaaaa      1140
ggccagcaaa aggccaggaa cggtaaaaag gcccgcgttgc tggcgcccccccttccataggctc      1200
cgccccccctg acgagcatca caaaaatcgat cttttttttttt tttttttttttt tttttttttttt      1260
ggactataaa gataccaggc gtttccccctt ggaagtcggat tttttttttttt tttttttttttt      1320
accctggccca tttttttttttt tttttttttttt tttttttttttt tttttttttttt tttttttttttt      1380
catagctcac gctgttaggtt tttttttttttt tttttttttttt tttttttttttt tttttttttttt      1440

```

-continued

gtgcacgaac ccccggttca gccggaccgc tgccgcttat ccggtaacta tcgtcttgag
tccaaccgg taagacacga cttatcgcca ctggcagcag ccactggtaa caggattagc
agagcgaggat atgttagggg tgctcacagag ttcttgaagt ggtggctaa ctacggctac
actagaagga cagtatttg tatctgcgtc ctgctgaagc cagttacctt cgaaaaaga
gttggtagct cttgatccgg caaacaaaacc accgctggta gcgggtgttt tttgtttgc
aaggcagcaga ttacgcccag aaaaaaagga tctcaagaag atcccttgat cttttctacg
gggtctgacg ctcagtggaa cggaaaactca cgttaaggga ttttggtcat gagattatca
aaaaggatct tcacctagat ccttttaaat taaaaatgaa gttttaatc aatctaaagt
atatatgagt aaacttggtc tgacagttac caatgctta tcagtgggc acctatctca
gegatctgtc tatttcgttc atccatagtt gectgactcc ccgtctgtt gataactacg
atacgggagg gcttaccatc tgccccca gctgcaatga taccggcaga cccacgctca
ccggctccag atttatcagc aataaaccag ccagccggaa gggccgagcg cagaagtgg
cctgcaactt tatccgcctc catccagtct attaatttgtt gccgggaagc tagagtaagt
agttcggccag ttaatagttt ggcacacgtt gtggccattt ctacaggcat cgtgggtca
cgctctgtc ttggatggc ttcatcagc tccgggtttcc aacgatcaag gcgagttaca
tgatccccca tggatggc aaaaagcggtt agtccttcg gtcctccgat cggttcaga
agtaagttgg cccgcaatgtt atcaatcatg gttatggcag cactgcataa ttctcttact
gtcatgccat ccgtaaatgtt ctttctgtc actgggtgagt actcaacccaa gtcattctga
gaatagtgtt tgccggcacc gagttgtct tggccggcgt caatacgggaa taatccgc
ccacatagca gaactttaaa agtgctcatc attggaaaac gttcttcggg gcgaaaactc
tcaaggatct taccgtgtt gagatccagt tcgatgtaac ccactctgtc acccaactga
tcttcagcat cttttacttt caccagcggtt tctgggttag caaaaacagg aaggccaaaat
gcccggaaaaa aggaaataag ggccgacacgg aatgttgaa tactcatact cttccctttt
caatagctcc aaggcaacaa attgactact cagaccgaca ttcatcgtt attgatttt
aatcaacgat aaacggaaatg gttacttgaa tgatttactt ttatgatcat tgtttactaa
ttacctaataat aggattttat atggaaattgg aagaataagg gaaatttcag atgtctgaaa
aaggcggagga gggtaactaat cattcaagcc catttctgc cagtaattgc ttccataagct
tcaataacttttactt cttgatagca atttctgtc ccatggctac gccccttttgc
ccattcaatc cggtggccgtt caaccaatct ctgagaaact gcttacgtt actctcttgc
gatttaccca cttggtaagt cttttgatttccaa caaaatcttag aagaatctgg agttaaaact
tcatctacta gtaccaatttccaa attgttttcg tccagttccaa attcgaattt cgtatcagca
ataatgatcc ctttcaaaaag ggcgaagttt tttgcagcag aataacaactc gaccggcttgc
acagcgacct tctcacaaat gtctttacccat acaatctcg cagcttggttc aatagagatg
ttttcatcgat gttcacccgtt ttcagttccaa gttgaaggtt tgaaaatccgg agttggaaag
ggcgtcgatctt cttggtaagt cttggccgtt accttgcgtc catggacagt ttttgggtt
ttgtactctt tccatgcact tccagttccaa gttgaaggtt tgaaaatccgg agttggaaag
agtctgtgtt cttttactat caaggatctt ccctctaaat gagatgttgc ttttttttgc
gacagtttttgc atggtagttaa agcaaagact tccctgtcat tagaagcaac caaatgttcc
tttatgttagg gtcggaaaaa atcaaaccag aaaactgaga gctgagttca aatcttttttgc
3720

-continued

ttatcaggaa taccgttgt cataatcaca tcgtaagcg agataacgac agttgcgac	3840
aacagcaagt tggcttcata gactgcata atgtctctaa ccttccctt ggcgattaaa	3900
ggtaggattc cggtccagatc agtggtcaca atggacatac ttggaaggat acagcaaagt	3960
gtgttggaaag cgatgacaca tggaaaggaa ttttcgagt ttcctagagt agtatattgg	4020
ggcgggtggaaa gttcagatgt ttaatgccta atactcttat actcttcaa gcgcggaaagt	4080
gtttctgcca acctgacttt ttctgataa atgaatcggt caagtggagt atttaaacca	4140
tgattaagtt acgtgatttgc acactggata aggtcgaaaa atatccgtat tcataaacga	4200
ttatttggtaa aagttacaaa ataccactaa ttacggagaa gcttagtaac agttatcatc	4260
tcttggtcga ttaacgccta caattccat tcgcccattca ggctgcgca ctgtgggaa	4320
ggcgatcggttgcggggcc ttcgcttattca cgccaggggcc tcgaggacaca aacgaacgtc	4380
tcacttaatc ttctgtactc tgaagaggag tgggaaatac caagaaaaac atcaaactcg	4440
aatgatttc ccaaaccctt accacaagat attcatcgc tgcgagatag gctgatcagg	4500
agcaagctcg tacgagaaga aacaaaatga caaaaaaaaaat cctataactat atagttaca	4560
aataaaaaaaat tatcaaaaat gaagcctgca tctctcaggc aaatggatt ctgacatcct	4620
cttgatttata atctagcaag accgggtttc tcgtaagtgc ccaacttgaa ctgaggaaca	4680
gtcatgtcta aggctacaaa ctcacatgtatc atgatgtatc tggtcgacgg cgctattcag	4740
atcccttttctt gagatgagtt ttgttcggg cccaaaaata acttccaaat cttcagttat	4800
gtcaagcgat aggttaagaa tggtgatctt attatcatc tcataatgcg atctatcg	4860
caatgaactt tcttcgccccat ttttttgcattttaataactg tctttcaacgc ttatgttgc	4920
ttgttaatttta ttcttgcga taatgattttt ttcaatatttgc ttatccctca aagtatttgc	4980
gtgaatgcctt gtcaaattctt cggaaatatgc acttcgaacaa ttttttaagg tattttttc	5040
gaaaatgaac ttagtttctt cgtactcacc tctttggtag cccaaatgtt ctccatcatc	5100
aacataaaaga tctccaaacgg ctgcgtccctc ggtatcaggg gctataacta ttacatatgg	5160
atcggttttca attaacatttgc aagatcttctt atacttataatc ttcatatgtatc taatgtgacc	5220
gccttcaata aataatggaa ttttatccaa tggtgacaaatc atattctttt ctatcaaattc	5280
agtaccatttgc ttataaaatgc agtgtaaaga tgcgaatttca tagaatatac cgggtgggaa	5340
aaccatttcc gtttctgattt gaccaggctc cgtgacaggtt tgactaata gacctgaatttgc	5400
actccagttaa aattggttat cgtatgtataa caattcagca aattcaggggtt gttcaataaaaat	5460
cattggatttca attatcgaaatccatcgtgac acttggatttgc ttcatatgtatc taatgtgacc	5520
aggtagcagg aaatatctca atttggataat atcacgtactt atcgacttca aagggttcatc	5580
gaataagttat ggttcttcttcc tcttgggtgtc tataatggggcg tggctcttcaaaaatgggtt	5640
ccataagccc gcttgggtacc aacgtgcaat caattcaggtt gtaggatctt cagcaagcc	5700
agctatgtcg gcttcttcaaaatgcatacc agcaatgttgc tttggacaaatccatcgttca	5760
ggaatctt aagtaatccc aattggccac attgtcacca gtcacatgtgg cagcagttac	5820
tttgagggccg gcaaaaaaaaaatgcatacc gcttcttgc tggaaatggatcgg atgggttgc	5880
aatcgatttttatttgcgttca gtagtgcattc atgcactgtatc agaccatata tggatggac	5940
ggatcttcc tcaatgtatc tggctgtatc caaatcttttggatgttgc tctctggggcc	6000
atcgaaaatc gaaaggctgttca gtagtgcattc atgcactgtatc agaccatata tggatggac	6060
cagatccata aaccgttgcgaaaaggactt ccaaatcttttggccatatttgc tggatggac	6120
atcaatccat atagaatttgc tggccagca atgacccatcata tagtcatatcatttgc tggatggac	6180

-continued

cttactgct acatttcat taattaccct gtcactgatt tcataatctt tcttaaatg	6240
aggatcgatt agtacgcaca gatttctacc caacttttt aatttgata acagccttt	6300
tggattggaa aaggagtgtct gttccaagt aaaatattt ttgtcggtcg tatactccaa	6360
gtccaaccaa ataaaatcgta aaggaatcat atgagcatcc atctgagagt ccactgtgag	6420
aacgtccatc tcatcattat aattccatct acattgtatgg taccctatacg aggaaatggg	6480
cggtaaaaag ggtctaccag tcaaatcggt aaatttgta atgatagttt gaatatctgg	6540
ccccagggac atgactacat ctatgacacc atttcggag atccaatgag tcatacgaaaa	6600
attttactg gtgtcatact ttatgtctca ccaagtgtca gctgcattga cccaaaagat	6660
agatgtggac gaagatgaaa acatgaatgg gatcgaaccg tacattgggtt ggctggtacc	6720
gatgttgtac tcaaagacat caacgttgaa aagcctgttag ggtccttttc caccgtaaat	6780
gtccatcagc cttagcgacg tcgcattgttc cggtataccg tagacattag tagaaccat	6840
gaaagagaaa tctagcgca cccattcagg ccccaaaggc atagagtcat gcttgaata	6900
caagaaatttgc tccttaaaca tggtgaaatg tgtttcttgc ggcagcacgt gtgcgaagtt	6960
ttcctgctta gtttatgtat gttcaatgtt caggaaatgg tgctcggttta caataagttt	7020
cagcgcattt tgccagtaaa cttcaatttgc aaaagggttca gcaaagatttt ctacggatac	7080
atcaccgttt cgaagatgaa atgtgtctgc agtggagttt gaaagtgaca aaaatgaga	7140
tattttcgac cagaatgagt tcacagtttgc ttttgcgttta aggaagtggaa attgtggaa	7200
actggccttgc ttcgccttc cttgaaattt cttgtcgaaat gcgtacttcc aggtctcatt	7260
gaaccgttgtt gaagagatca acaaaccgct gctgttgggtt ggcattctct ctttctcatt	7320
tatagtgaac cttactgagt gatcctgttgc aaaagagaga gagaatggga actgaacggc	7380
tatatcatcg ccctccaatc ttggatatacg ttttattatg gtgcattgaa gcacatttc	7440
taaaggatcg tgcataatcg actcgccgtc cactttgttgc tagcgtatgat gagattggc	7500
aatatttcttgcataaaatcc ttgttctatg gcaaaaacca gattgcgcac acttctttaa	7560
tagatagtcg gtaaacgcattt gcgaaaaaggc ggtaaagaag accaattggc atacgagccaa	7620
tttcaaaaagg accatctcg	7639

<210> SEQ ID NO 22
<211> LENGTH: 5742
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pYPTIZAGLSII.

<400> SEQUENCE: 22

cgaacaaaaa ctcatctcg aagaggatct gaatagcgcc gtcgaccatc atcatcatca	60
tcattgagtt tgtagcctta gacatgactt ttccctcgat caagttgggc acttaacgaga	120
agaccggctt tgcttagattc taatcaagag gatgtcagaa tgccatttgc ctgagagatg	180
caggcttcatttttgcataact tttttatttgc taaccttatat agtataaggat ttttttgtc	240
attttgcattttc ttctcgatcg agcttgcgtcc tgatcagcctt atctcgccgc tgatgaat	300
cttgcgttag gggtttggaa aaatcattcg agtttgcgtt ttttcttgcgtt atttccact	360
ccttttcaga gtacagaaga ttaagtgaga ctttcgttgc tgccgttccccc ccacacacca	420
tagcttcggaa atgtttctac tccttttttgc ctcttcggaa ttttcttgcgtt ctccgcgtt	480
cggcgatccatc cttcaaaaaca cccaaaggcaca gcatactaaa ttttcccttc ttcttcgtt	540

-continued

agggtgtcgt taattacccg tactaaagggt ttggaaaaga aaaaagagac cgcctcggtt 600
 cttttccttc gtcggaaaag gcaataaaaaa ttttatcac gtttctttt cttgaaattt 660
 ttttttttag ttttttctc tttcagtgac ctccattgtat atttaagtta ataaacggtc 720
 tc当地tct caagtttcag tttcattttt cttgttctat tacaactttt tttacttctt 780
 gttcattaga aagaaagcat agcaatctaa tctaaggggc ggtgttgaca attaatcatc 840
 ggcatacgat atcggccatag tataatacgca caaggtgagg aactaaacca tggccaagtt 900
 gaccagtgcg gttccgggtgc tcaccggcg cgacgtcgcc ggagcggtcg agttctggac 960
 cgaccggctc gggttctccc gggacttcgt ggaggacgac ttccgggtcg tggccggga 1020
 cgacgtgacc ctgttcatca ggcgggtcca ggaccagggtg gtggccgaca acaccctggc 1080
 ctgggtgtgg gtgcggccgc tggacgagct gtacggcgag tggcggagg tcgtgtccac 1140
 gaacttccgg gacgcctccg ggccggccat gaccgagatc ggcgagcagc cgtggggcg 1200
 ggagttcgcc ctgcgcgacc cggccggcaaa ctgcgtgcac ttccgtggccg aggagoagga 1260
 ctgacacgtc cgacggcgcc ccacgggtcc caggcctcg agatccgtcc ccctttcct 1320
 ttgtcgatat catgtatcattt gttatgtcact gtttacatc acggccctccc cccacatccg 1380
 ctcttaaccga aaaggaaagga gtttagacaac ctgaagtctt ggtccctatt tatttttta 1440
 tagttatgtt agtattaaga acgttattttt tatttcaaattt ttttctttt tttctgtaca 1500
 gacgcgtgta cgcatgtaac attatactga aaaccttgcgt tgagaagggtt ttggggacgct 1560
 cgaaggctt aatttgcaag ctggagacca acatgtgagc aaaaggccag caaaaggccca 1620
 ggaaccgtaa aaaggcccgcg ttgctggcg tttccatag gctccggccc cctgacgagc 1680
 atcacaaaaaa tcgacgctca agtcagagggt ggcaaaaccc gacaggacta taaagatacc 1740
 aggegttcc cccttggaaacgc tccctcggtgc gctctctgt tccgaccctg ccgttaccg 1800
 gatacctgtc cgccttctc cttccggaa gctggcgct ttcctcaatgc tcacgtgta 1860
 ggtatctcg ttccgtgttag gtcgttcgct ccaagctggg ctgtgtgcac gaaccccccg 1920
 ttcagcccgaa ccgcgtgcgc ttatccggta actatcgctc tgagtccaaac ccggtaagac 1980
 acgacttatac gccactggca gcagccactg gtaacaggat tagcagagcg aggtatgttag 2040
 gcggtgtac aggttcttg aagtgggtggc ctaactacgg ctacactaga aggacagtt 2100
 ttggtatctg cgctctgtc aagccagtta cttccggaaa aagagtttgtt agctttgtat 2160
 cccggcaaaaca aaccaccgctt ggttagccggt gttttttgtt ttgcggacgac cagattacgc 2220
 gcagaaaaaaa aggatctcaa gaagatctt tgcgttttacgggttctt gacgctcgt 2280
 ggaacgaaaaa ctcacgttaa gggatccggt tcatgagatc agatctatga tgagtccaa 2340
 tctgttttcca cagacgagta caaggacagg caaaaggaaat tggaaagaatg tgctaaaccc 2400
 ataatgagca agttctatgg agtcgtgttgg ggagctccgt gtcgttgcac ttggggcttc 2460
 cctggagggtt tccctggcgcc agctggcgca gctggcggtt ccccgagggtt tgctggccca 2520
 ggcggagaca ggcggaccaac cgtggaaagaa gtcgattaag caattcaacg gataaattct 2580
 ggttaataataataacgtga ataggaaattt aaggaaattt tggatctaat aatgtgtgt 2640
 atgcccacat cggggcatgtt agattgtata gtcgtgtca cactataata agccagccaa 2700
 aaccctttaaa ccagttggcc tccactaattt agtgtactac ccaatcttc ctcttcgggt 2760
 gtcttttata aggacagatt cacaagctct tggccggccaa tacacacata cacacagaga 2820
 taatagcgtt cgaatttcacg tggcccgcc ggcgttccgt gatcggtacc tcgagatgg 2880

-continued

cctttgaaa tggctcgat gccaattggt cttctttacc gcttttcgc atgcgttac	2940
cgactatcta ttaaaagaagt gtgcgcatac tgggtttgc catagaaaca gggtttatgc	3000
agaaaatatt gccaaatctc atcactgcta ttacaaagtg gacgcgagt ctattgcaca	3060
cgatcctta gagaatgtgc ttcatgtac cataattaaa actataccaa gattggaggg	3120
cgatgatata gccgttcagt tcccattctc tctctttt ttacaggatc actcagtaag	3180
gttcactata aatgagaaaag agagaatgcc aaccaacagc agcggtttgt tgatctttc	3240
acaacggttc aatgagacct ggaagtacgc attcgacaag aaattcaag aggaggcgaa	3300
caggaccagt atccacaat tccacttcc ttaagcaaaaa caaactgtga actcattctg	3360
gtcgaaaata ttttcatttt tgcactttc aaactccact gcagacacat ttcatctcg	3420
aaacggtgat gtatccgtag aaatcttgc tgaaccttt caattgaaag ttactggca	3480
aatgcgcgtg aaacttattt taaacgagca aaatttcctg aacattgaac atcatagaac	3540
taagcaggaa aacttcgcac acgtgctgcc agaagaaaaca actttcaaca tgtttaagga	3600
caatttcctg tattcaaagc atgactctat gcctttgggg cctgaatcgg ttgcgttaga	3660
tttcttttca atgggttcta ctaatgtcta cggatacccg gaacatgcga cgtcgctaag	3720
gctgatggac acttcagggtg gaaaggaacc ctacaggctt ttcaacgttg atgtcttga	3780
gtacaacatc ggtaccagcc aaccaatgtc cggttcgatc ccattcatgt tttcatcttc	3840
gtccacatct atcttttggg tcaatgcgc tgacacttgg gttagacataa agtatgacac	3900
cagtaaaaat aaaacgatga ctcattggat ctccgaaaat ggtgtcatag atgtgtcat	3960
gtccctgggg ccagatattc caactatcat tgacaaattt accgatttga ctggtagacc	4020
cttttaccg cccatttcc ctagggta ccatcaatgt agatgaaatt ataatgtga	4080
gtatggacgtt ctcacagtgg actctcagat ggatgctcat atgattcctt acgattttat	4140
ttgggtggac ttggaggtata cgaacgacaa aaaatattt acttggaaagc agcactcctt	4200
tcccaatcca aaaaggctgt tatccaaatt aaaaaagtgg ggttagaaatc ttgtcgact	4260
aatcgatcct cattaaaga aagattatga aatcagtgc acggtaatta atgaaaatgt	4320
agcagtcaag gatcacaatg gaaatgacta tgcgttgc tgcgtggccag gtaattctat	4380
atggattgtat accataagca aatatggcca aaagattttgg aagtccttt tcgaacgggtt	4440
tatggatctg cccgctgatt taactaattt attcatttgg aatgatatga acgagcttc	4500
gattttcgat ggcccaagaga ccacagctcc aaaagatttgc attcacgaca attacattga	4560
ggaaagatcc gtccataaca tatatggct atcagtgcat gaagctactt acgacgcaat	4620
aaaatcgatt tattcaccat ccgataagcg tctttccctt ctaacaaggg cttttttgc	4680
cggctctcaa cgtactgtg ccacatggac tggtgacaaat gtggccaatt gggattactt	4740
aaagatttcc atccatgtgg ttctgtcaaa caacattgtt ggtatgcac ttataggagc	4800
cgacatagct ggctttgtg aggatctac acctgaatttgc attgcacgtt ggtaccaagc	4860
gggcttatgg tacccatgtt ttagagcaca cgcccatata gacaccaaga gaagagaacc	4920
atacttattc aatgaacctt tgaagtcgtat agtacgtgtat attatccat tgagatattt	4980
cctgctaccc accttataca ccatgttca taaatcaagt gtcactggat ttccgataat	5040
gaatccatg tttattgaac accctgaatt tgctgaatttgc tatcatatcg ataaccaatt	5100
ttactggagt aattcagggtc tattgtcaaa acctgtcaccg gagcctggtc aatcagaac	5160
ggaaatgggtt ttcccacccg gtatattcta tgaattgcac tctttacact ctttataaa	5220
caatggtaact gatttgatag aaaagaatat ttctgcacca ttggataaaa ttccatttt	5280

-continued

tattgaaggc ggtcacatta tcactatgaa agataagtat agaagatctt caatgttaat	5340
aaaaaacgat ccatatgtaa tagttatagc ccctgatacc gagggacgag ccgttgaga	5400
tctttatgtt gatgatggag aaactttgg ctaccaaaga ggtgagtacg tagaaactca	5460
gttcatttgc gaaaacaata ccttaaaaaa tgttcgaagt catattccg agaatttgac	5520
aggcattcac cacaatactt tgaggaatac caatattgaa aaaatcatta tcgcaaagaa	5580
taatttacaa cacaacataa cgttgaaaga cagtattaaa gtcaaaaaaa atggcgaaga	5640
aagttcattg ccgacttagat cgtnatgaa gaatgataat aagatcacca ttcttaacct	5700
atcgcttgac ataactgaag attgggaagt tattttggg cc	5742

<210> SEQ ID NO 23
<211> LENGTH: 7256
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pYPT1ADE1glsII.

<400> SEQUENCE: 23

gtcgaccggc tgcattaatg aatcgccaa cgcgccccca gaggcggttt gcgtattggg	60
cgccttccg cttccctcgct cactgactcg ctgcgcctcg tcgttcggct gcggcgagcg	120
gtatcagctc actcaaaggc ggtataacgg ttatccacag aatcaggggtaa acgcaggg	180
aagaacatgt gagcaaaaagg ccagcaaaa gccagggacc gtaaaaaggc cgcttgctg	240
gcgttttcc ataggctcg ccccccgtac gagcatcaca aaaatcgacg ctcaagtctc	300
aggtggcgaa acccgacagg actataaaga taccaggcgt ttccccctgg aagctccctc	360
gtgcgcctc ctgttccgac cctgcccgtt accggatacc tgtccgcctt tctcccttcg	420
ggaagcgtgg cgcttctca tagtcacgc tgcgttgcgt tcaagttcggt gtaggtcggt	480
cgctccaagc tggcgtgtgt gcacgaacc cccgttcagc ccgaccgtcg cgccatatcc	540
gttaactatac gtcttgcgtt caacccggta agacacgact tatcgccact ggcagcagcc	600
actggtaaca ggatttagcag agcgaggtat gtagggcgtt ctacagatcc ttgttgttgg	660
tggcttaact acggctacac tagaaggaca gtatggta tctgcgtct gctgaaggcca	720
gttacccctcg gaaaaagagt tggtagctc tgatccggca aacaaaccac cgctggtagc	780
ggtggtttt ttgttgttgcgaa gcagcagatt acgcgcagaa aaaaaggatc tcaagaagat	840
cctttgatct tttctacggg gtctgacgct cagtgaaacg aaaactcagc ttaagggatt	900
ttggcgtatgtt gattatcaaa aaggatctt acctagatcc ttttaaattttaaaaatgtt	960
ttaaatcaatctaaatgtt atatgagttaa acttggcgtt acagttacca atgtttatc	1020
agtggcgttacac cttatctcagc gatctgttata tttcggttcat ccatacgatgc ctgactcccc	1080
gtcggtttaga taactacgtt acggggggcc ttaccatctg gccccagtc tgcaatgtata	1140
ccggcggacc caccgttccacc gggtcccgat ttatcgacaa taaaccaggcc agccggaaagg	1200
ggccggccgtca gaagtggtcc tgcacatttca tccgccttca tccagtttat taattttgc	1260
cggggagctttaa gagtaagttag ttccgttgcgtt aatagttgc gcaacgttgc tgccattgt	1320
acaggcatcg tgggtgttacg ctcgtcggtt ggtatggctt cattcagtc cgggtcccaa	1380
cgatcaaggc gagttacatg atccccatg ttgtgcaaaa aagcggttag ctcccttcggt	1440
cctccgtatcg ttgtcagaag taagttggcc gcaatgtttat cactcatgtt tatggcagca	1500
ctgcataatt ctcttactgtt catgcccatttca gtaatgttgc ttctgttgc tggtgttgc	1560

-continued

tcaaccaagt cattctgaga atagtgtatg cgccgcaccga gttgctttg cccggcgtca 1620
 atacgggata ataccgcgcc acatagcaga actttaaaag tgctcatcat tggaaaacgt 1680
 tcttcgggac gaaaactctc aaggatctt ccgctgtta gatccagttc gatgtAACCC 1740
 actcggtcac ccaactgtatc tttagtcttca ccagcgttcc tgggtgagca 1800
 aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg cgacacggaa atgttgaata 1860
 ctctactct tccttttca atagctccaa ggcaacaat tgactactca gaccgacatt 1920
 cattcggtat tgatTTTaaa tcaacgataa acggaatggt tacttgaatg atttacttt 1980
 atgatcattt tttactaatt acctaataag gatTTTatat ggaattggaa gaataaggaa 2040
 aatttcagat gtctgaaaaa ggccgaggagg gtactaatca ttcaagccca ttcttgcca 2100
 gtaattgctt cataagcttc aataacttt tctttactct tgatagoaat ttctgoatcc 2160
 atggctacgc cctctttgcc attcaatccg ttggccgtca accaatctct gagaaactgc 2220
 ttatcgtaac tctcttgca tttacccact tggtaagtct tttgattcca aaatctagaa 2280
 gaatctggag taaaacttc atctacttagt accaattcat tgTTTcgtc cagtcoaat 2340
 tcgaatttcg ttcagcaat aatgatcccc ttcaaaaaggg cgaagTTTT tgccggagaa 2400
 tacaactcga ccgccttgac agogacccctc tcacaaatgt cttaacctac aatctcagca 2460
 gtttgtcaa tagagatgtt ttcatcggt tcacccctgtt cagcttcgt tgaagggttg 2520
 aaaatcggag ttggaaaggc gtgcgtctct tgaaggttct cgtttcaac cttgactcca 2580
 tggacagttt ttgagttttt gtactcttc catgcacttc cagtgtatgta acctctgaca 2640
 atggcttcca aaggtatcag tctgtgttt tttactatca aggatcgatcc ctctaattga 2700
 gatttgtatt ttcttcaga cagTTTTgtt ggttagtaaag caaagacttc cttgtcatta 2760
 gaagcaacca aatgatttt tatgttagggt gccaaaaat caaaccagaa aactgagagc 2820
 tgagtcaaaa tctttccctt atcaggaata ccgtttgtca taatcacatc gtaaggggag 2880
 atacggtcag ttgcgacgaa cagcaagttt ttctcatcga ctgcataat gtctctaacc 2940
 ttcccttgg cgattaaagg taggatccc tccagatcag tgTTTcacaat ggacatactt 3000
 ggaaggatac agcaaagtgt gttggaaagcg atgacacatg gaaaggatt ttgcagttt 3060
 cctagagtag tataattgggg cggtgaaagt tcagatgtt aatgcttaat actcttatac 3120
 tcttcaaagc gcccagtgt ttctgccaac ctgactttt tctgaataat gaatcggtca 3180
 agtggaggtat ttaaaccatg attaagtatc gtgatttgcg actggataag gtcggaaaat 3240
 atccgtattt ataaacgatt atggtaaaa gttacaaat accactaatt acggagaagc 3300
 ttagtaacag ttatcatctc ttggatcgatt aacgcttaca atttccattc gccattcagg 3360
 ctgcgcact gttggaaagg gcatcggtt cgggccttt cgctattacg ccagggcctc 3420
 gaggcacaaa cgaacgtctc acttaatctt ctgtactctg aagaggagtgg gaaaatacc 3480
 agaaaaacat caaactcgaat tgatTTTccc aaaccctac cacaagatat tcatcagctg 3540
 cgagataggc tgatcaggag caagctcgta cgagaagaaa caaaatgaca aaaaaaatcc 3600
 tataactat atggtaaaa taaaaaaatgta taaaaatgta agcctgoatc tctcaggcaa 3660
 atggcattct gacatcctct tgatttagat ctgcggatc cggtcttcgt gtaagtggccc 3720
 aacttgaact gaggaacagt catgtctaa gctacaaact caatgtatgat gatgtatgt 3780
 gtcgacggcg ctattcagat cctcttgcgat gatgagttt tggatggccc caaaaataac 3840
 ttcccaatct tcatgttgcgat caagcgatag gtttggatgatcattat tattttttttttt 3900

-continued

atatgacgat	ctagtccggca	atgaacttgc	tgcgcatttt	tttttgactt	taatactgtc	3960
tttcaacgtt	atgttgtgtt	gtaaattattt	ctttgcgata	atgatttttt	caatattgg	4020
atttcctcaaa	gtattgtgg	aatgtcgt	caaattctcg	ggaatatgac	tgcgaacatt	4080
tttaaggta	ttgttttgcg	aaatgaactg	agtttctacg	tactcacctc	tttggtagcc	4140
aaaagtttct	ccatcatcaa	cataaagatc	tccaaacggct	cgtccctcg	tatcaggggc	4200
tataactatt	acatatggat	cgtttttcat	taacattgaa	gatcttctat	acttatctt	4260
catagtgata	atgtgaccgc	cttcaataaa	taatgaaatt	ttatccaatg	gtgcagaaat	4320
attcttttct	atcaaatcag	taccattgtt	tataaaagag	tgtaaagatg	cgaattcata	4380
gaatataccg	ggtggggaaaa	ccatttcgt	ttctgattga	ccaggctccg	tgacaggttt	4440
gactaataga	cctgaattac	tccagtaaaa	ttggttatcg	atatgataca	attcagcaaa	4500
ttcagggtgt	tcaataaaca	ttggattcat	tatcgaaat	ccagtgcac	ttgatttatg	4560
aaacatggtg	tataaggtag	gtagcaggaa	atatctcaat	tggataat	cacgtactat	4620
cgacttc当地	ggttcattga	ataagtatgg	ttctcttc	ttgggtctca	tatggcg	4680
tgctctaaaa	aatgggtacc	ataagccccg	ttggtagcaa	cgtgcaatca	attcaggtgt	4740
aggatcctca	gcaaagccag	ctatgtccgc	tcctataaaat	ggcataccag	caatgttgc	4800
tgacagaacc	ataggaatgg	aaatctttaa	gtaatccaa	ttggccacat	tgtcaccagt	4860
ccatgtggca	gcagtacgtt	gagagccggc	aaaaaaagcc	cttggtagaa	ggaaaggacg	4920
cttatcgat	ggtgaataaa	tcgattttat	tgcgtcgtaa	gtagcttcat	gcactgatag	4980
accatatacg	ttatggacgg	atctttctc	aatgtatgg	tcgtgaatca	aatctttgg	5040
agctgtggtc	tctggccat	cgaaaatcga	aggctcg	atatcattcc	aaatgaataa	5100
attagttaaa	tcagccggca	gatccataaa	ccgttgcgaa	aaggactcc	aatcttttg	5160
gccatatttgc	tttatggat	caatccat	agaattac	ggccagcaat	gacctacata	5220
gtcatttcca	tttgatct	tgactgat	atttcattta	attaccctgt	cactgatttgc	5280
ataatcttgc	tttaaatgag	gatcgattag	tacgacaaga	tttctacca	actttttaa	5340
tttggataac	agccttttg	gattggggaa	ggagtgctc	ttccaagtaa	aatattttt	5400
gtcggtcgta	tactccaatg	ccaaaccaat	aaaatcgtaa	ggaatcatat	gagcatccat	5460
ctgagagtcc	actgtgagaa	cgtccatctc	atcattataa	ttccatctac	attgatggta	5520
ccctatagag	gaaatggcg	gtaaaaagg	tctaccagtc	aaatcgtaa	atttgcata	5580
gatagttgga	atatctggcc	ccagggacat	gactacatct	atgacaccat	tttcggagat	5640
ccaatgagtc	atcgtttat	tttactgg	gtcatactt	atgtctaccc	aagtgtcagc	5700
tgcattgacc	caaaagatag	atgtggacg	agatgaaaac	atgaatggg	tgcgaaaccgt	5760
cattgggtgg	ctggtaccga	tgttgtactc	aaagacatca	acgttggaaa	gcctgttaggg	5820
ttcccttcca	cctgaagtgt	ccatcagect	tagcgacgt	gcatgttccg	gtataaccgt	5880
gacatttagta	gaacccatga	aagagaaatc	tagcgacacc	gattcaggcc	ccaaaggcat	5940
agagtcatgc	tttgaataca	agaaattgtc	cttaaacatg	ttgaaagttg	tttcttctgg	6000
cagcacgtgt	gcgaagtttt	cctgcttagt	tctatgtatgt	tcaatgtca	ggaaatttt	6060
ctcgtttaca	ataagttca	gcgcattttg	ccagtaact	ttcaattgaa	aaggttcagc	6120
aaagatttct	acggatacat	caccgttgc	aaagatgaaat	gtgtctgcag	tggagttga	6180
aagtgcacaaa	aatgaagata	tttgcacca	aatgagtc	acagttgtt	tttgcttaag	6240
gaagtggaaat	tgtggaaatac	ttgtccctgtt	cgccctctct	tgaaattct	tgtcgaatgc	6300

-continued

gtacttccag gtctcattga accgttgtga agagatcaac aaaccgcgtgc tgttgggtgg 6360
 cattctctt ctctcattta tagtgaacct tactgagtga tcctgtaaaa aagagagaga 6420
 gaatggAAC tgaacggcta tatcatcgcc ctccaatctt ggtatagttt taattatgg 6480
 agcatgaagc acattctcta aaggatcgta tgcaatagac tcggcggtcca ctttgtata 6540
 gcagtgtga gatttggcaa tattttctgc ataaaccctg tttctatggc aaaaccaga 6600
 ttgcgcacac ttctttaata gatagtcggt aaacgcgtgc gaaaaagcg 6660
 caattggcat acgagccatt tcaaaaggac catctcgagg taccgatccg agacggccgg 6720
 ctggggcacg tgaatttcac tgctattatc tctgtgtgtta tttgtgtatt gggcaacaag 6780
 agcttgtgaa tctgtcctta taaaagacac ccgaagaggc aagattgggt agtacactaa 6840
 ttagtggagg gcaactgggt taggggtttt ggctggctta ttatagtgta agcgatacta 6900
 tacaatctac gatgcccgtat gtccggcatac agcacattat tagatccaaa atttccttaa 6960
 ttccatttc acgttatata tattaaccag aatttatccg ttgaatttgct taatcgactt 7020
 ctccccacggt tggtcccgctg tctccgcotg gggcagcacc acctggggca ccgcacgtg 7080
 cgccagctcc gccaggaaa cctccaggaa agccaccagg agctccacca ggagctccac 7140
 cagcagctcc atagaacttg ctcaattattt ggttagcaac ttcttccaat ttctttgcc 7200
 tgccttgta ctgcgtgtg gaagcagatt gtgactcatc atagatctga tctcat 7256

<210> SEQ ID NO 24
 <211> LENGTH: 5721
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pGAPZAglsIIHDEL.

<400> SEQUENCE: 24

tcgagatggt cttttgaaa tggctcgat gccaatttgtt cttcttacc gcttttcgc 60
 atgcgtttac cgactatcta taaaagaagt gtgcgcatac tgggtttgc catagaaaca 120
 gggtttatgc agaaaatatt gccaaatctc atcaactgcta ttacaaatgt gacgcccagt 180
 ctattgcaca cgatcctta gagaatgtgc ttcatgtcac cataattaaa actataccaa 240
 gattggaggg cgatgatata gccgttca gtcatttc tctctttttt ttacaggatc 300
 actcaactaa gttcaactata aatgagaaag agagaatgcc aaccaacagc agcggttgt 360
 tgcgttc acaacgggtt aatgagaccc ggaagtacgc attcgacaag aaatttcaag 420
 aggaggcgaa caggaccagt attccacaat tccacttc taaacgatc aaaaactgtga 480
 actcatttcg tgcggaaaata ttttcatttt tgcgttc aactccact gcagacacat 540
 ttcatcttcg aaacgggtat gtatccgtat aaatcttgc tgaacctttt caattgaaag 600
 ttactggca aatgcgtgc aacttattt taaacgatc aaatttcctg aacatgtga 660
 atcatagaac taacggaa aacttcgcac acgtgtgtcc agaagaaca acttcaaca 720
 tggtaagga caatttcctt tattcaaaatc atgactctat gcctttgggg cctgtatcg 780
 ttgcgttgc ttttcatttc atgggttctt ctaatgtctt cggatccacg gaacatgcga 840
 cgtcgctaa gctgtatggac acttcaggatg gaaaggaaacc ctacaggctt ttcaacgttg 900
 atgtcttgc tgcgtatccatc ggttccatc aaccaatgtt cggatccatc ccatttcatgt 960
 ttcatcttc gtcacatctt atcttttggg tcaatgtatc tgacacttgg gtagacataa 1020
 agtacatgtatc cagtaaaaat aaccaatgtt ctcattggat ctccgaaaat ggtgtatgt 1080

-continued

-continued

acccgtacta aaggtttggaaagaaaaaa gagaccgcctcgtttcttttcttcgtcga	3480
aaaaggcaat aaaatttttt atcacgtttcttttcttga aatttttttttttagttttt	3540
ttctcttca gtgacctcca ttgatattta agttaataaa cggtctcaa tttctcaagt	3600
ttcagttca ttttcttgt tctattacaa cttttttac ttcttgtca ttagaaagaa	3660
agcatagcaa tctaatactaa gggcgggtgt gacaattaat catggcata gtatatccgc	3720
atagtataat acgacaaggt gaggaaactaa accatggcca agttgaccag tgccgtccg	3780
gtgctcaccg cgccgcacgt cgccggagcg gtcgagttct ggaccgaccg gtcgggttc	3840
tccgggact tcgtggagga cgacttcgcg ggtgtggtcc gggacgacgt gaccctgttc	3900
atcagcgcgg tccaggacca ggtggtgcgg gacaacaccc tggcctgggt gtgggtgcgc	3960
ggcctggacg agctgtacgc cgagtggtcg gaggtcgtgt ccacgaacct ccgggacgcc	4020
tccggggccgg ccatgaccga gatcggcggag cagccgtggg ggcgggagtt cgccctgcgc	4080
gaccggccgg gcaactgcgt gcacttcgtg gccgaggagc aggactgaca cgtccgacgg	4140
cggccccacgg gtcggcggcc tcggagatcc gtcggcccttt tcggcgttcg atatcatgtt	4200
attagttatg tcacgcttac attcacgccc tccccccaca tcggctctaa ccgaaaagga	4260
aggagttaga caacctgaag tctaggtccc tatttttttttttataatgtt tgtagtatt	4320
aagaacgtta ttatatttc aaatttttct tttttttctg tacagacgcgtgtacgcgt	4380
taacattata ctgaaaacct tgcttgagaa ggttttggga cgctcgaagg cttaatttg	4440
caagctggag accaacatgt gagcaaaagg ccagcaaaagg gccaggaacc gtaaaaaggc	4500
cgcgttgcgt gcgcccccc ataggctccg ccccccgtac gagcatcaca aaaatcgacg	4560
ctcaagttag aggtggcgaa acccgacagg actataaaga taccaggcgt ttcccccgtt	4620
aagctccctc gtgcgtctc ctgttccgac cctggccgtt accggataacc tgcggccctt	4680
tctcccttcg ggaagcgtgg cgcttctca atgctcacgc tgtaggtatc tcagttcggt	4740
gtaggtcggt cgctccaagc tggcgtgtgt gcacgaaccc cccgttcagc ccgaccgctg	4800
cgccttatcc ggtaactatc gtcttgagtc caacccggta agacacgact tatcgccact	4860
ggcagcagcc actggtaaca ggatttagcaag agcgaggtat gtggcgggtg ctacagagtt	4920
cttgaagtgg tggcctaact acggctcacac tagaaggaca gtatttgta tctgcgtct	4980
gctgaagcca gttaccttcg gaaaaagagt tgtagctct tgatccggca aacaaaccac	5040
cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc	5100
tcaagaagat ctttgcgtt tttctacggg gtctgacgt cagtggaaacg aaaactcacg	5160
ttaagggatt ttggcatgc atgagatcag atcttttttgc tagaaatgtc ttgggtgcct	5220
cgtccaatca ggtagccatc tctgaaatat ctggctccgt tgcaactccg aacgcacgtc	5280
tggcaacgta aaattctccg gggtaaaact taaatgtgg gtaatggaa cagaaacgtc	5340
tctcccttc tctctccctc caccggccgt taccgtccctt aggaaattttt actctgtcg	5400
agagcttctt ctacggcccc ctggcagca tgctctccc agcattacgt tgccggtaaa	5460
acggaggtcg tgtacccgac ctgcgcaccc agggatggaa aagtccggc cgtcgtggc	5520
aataatagcg ggcggacgca tgcatgaga ttattggaaa ccaccagaat cgaatataaa	5580
aggcgaacac ctttcccaat tttggtttctt cctgacccaa agactttaaa ttatattat	5640
ttgtccctat ttcatcaat tgaacaacta ttgcggaaacg aggaatttcac gtggccacgc	5700
cggccgtctc ggatcggtac c	5721

-continued

<210> SEQ ID NO 25
 <211> LENGTH: 7230
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Sequence of plasmid pGAPADE1glsIIHDEL.

<400> SEQUENCE: 25

cgtactcacc	tctttggtag	ccaaaagttt	ctccatcatc	aacataaaga	tctccaacgg	60
ctcgccctc	ggtatcaggg	gctataacta	ttacatatgg	atcgaaaaatc	attaacattt	120
aagatcttct	atacttatct	ttcatatgtga	taatgtgacc	gccttcataa	aataatggaa	180
tttatccaa	tggtgcagaa	atattctttt	ctatcaaatac	agtaccattt	tttataaaag	240
agtgtaaaga	tgcgaaattca	tagaatatac	cgggtggaa	aaccattttc	gtttctgatt	300
gaccaggctc	cgtgacaggt	ttgactaata	gacctgaatt	actccagtaa	aattggttat	360
cgatatgata	caattcagca	aattcagggt	gttcaataaa	cattggattt	attatcgaa	420
atccagtgac	acttgattt	tgaatcatgg	tgtataaggt	aggtagcagg	aaatatctca	480
attggataat	atcacgtact	atcgacttca	aagggttatt	gaataagttat	ggttcttctt	540
tcttgggtgc	tatatggcg	tgtgctctaa	aaaatgggtt	ccataagccc	gtttggtacc	600
aacgtgcaat	caattcagggt	gttaggatctt	cagcaaaggcc	agctatgtcg	gtccctataa	660
atggcataacc	agcaatgtt	tttgacagaa	ccataggaat	ggaaatcttt	aagtaatccc	720
aattggccac	attgtcacca	gtccatgtgg	cagcgttacg	ttgagagccg	gaaaaaaaaag	780
cccttggtag	aaggaaagga	cgcttacgg	atggtaata	aatcgatttt	attgcgtcgt	840
aagtagctt	atgcactgat	agaccatata	tgttatggac	ggatcttcc	tcaatgtaat	900
tgtcgtgaat	caaatttttt	ggagctgtgg	tctctggccc	atcgaaaatc	gaaggctcg	960
tcatatcatt	ccaaatgaat	aaatttagtta	aatcagccgg	cagatccata	aaccgttcga	1020
aaaaggactt	ccaaatcttt	tggccatatt	tgttatgggt	atcaatccat	atagaattac	1080
ctggccagca	atgacctaca	tagtcattt	cattgtgatc	cttgcactgt	acattttcat	1140
taattaccct	gtcactgatt	tcataatctt	tctttaaatg	aggatcgatt	agtacgacaa	1200
gatttctacc	caactttttt	aatttgata	acagcctttt	tggattggaa	aaggagtgt	1260
gcttccaagt	aaaatatttt	ttgtcggtcg	tataactccaa	gtccaaacca	ataaaatcg	1320
aaggaatcat	atgagcatcc	atctgagagt	ccactgtgag	aacgtccatc	tcatcattat	1380
aattccatct	acattgtatgg	taccctatag	aggaaaatggg	cggtaaaaaag	ggtcttaccag	1440
tcaaatacggt	aaatttgtca	atgatagttt	gaatatctgg	ccccaggagac	atgactacat	1500
ctatgacacc	atttcggag	atccaatgag	tcatcgaaaaat	atttttactg	gtgtcataact	1560
ttatgtctac	ccaaatgtca	gctgcattt	ccaaaaagat	agatgtggac	gaagatgaaa	1620
acatgaatgg	gatcgaaacgg	tacattgggt	ggctggtacc	gatgttgc	tcaaagacat	1680
caacgttcaa	aaggctgttag	ggttccctt	cacctgaagt	gtccatcagc	cttagcgacg	1740
tcgcatgttc	cggatcaccg	tagacattag	tagaaccat	gaaagagaaa	tctagcgca	1800
ccgatttcagg	ccccaaaggc	atagagtcat	gtttgaata	caagaaattt	tccttaaaca	1860
tgttggaaatgt	tgtttcttctt	ggcagcacgt	gtgcgaagtt	ttcctgttta	gttctatgtat	1920
gttcaatgttt	caggaaatttt	tgotcgat	caataagttt	cagcgcattt	tgccagtaaa	1980
ctttcaattt	aaaaggttca	gcaaaagattt	ctacggatc	atcaccgttt	cgaagatgaa	2040
atgtgtctgc	agtggagttt	gaaaatgtaca	aaaatgtaga	tattttcgac	cagaatgtat	2100

-continued

tcacagttt ttttgctta aggaagtggaa attgtggaat actggtcctg ttgcctcct	2160
cttggaaattt cttgtcgaaat gcgtaacttcc aggtctcatt gaaccgttgtt gaagagatca	2220
acaaacccgt gctgttggtt ggcattctct ctttctcatt tatagtgaac cttaactgagt	2280
gatcctgtaa aaaagagaga gagaatggga actgaacggc tatatcatcg ccctccaatc	2340
ttaggtatagt ttaaattatg gtagcatgaa gcacattctc taaaggatcg tgtgcaatag	2400
actcgccgtc cactttgtaa tagcagtgtat gagatgttgc aatattttct gcataaaaccc	2460
tgtttctatg gcaaaaccca gattgcgcac acttctttaa tagatgtcg gtaaaacgcatt	2520
gcggaaaaagc ggttaaagaag accaattggc atacgagcca tttcaaaagg accatctcgat	2580
ggtaccgatc cgagacggcc gggtgggcca cgtgaattcc tcgtttcgaa atagttgttc	2640
aattgattga aataggggaca aataaaattaa atttaaagtc tttgggtcag gagaacccaa	2700
aattgggaaa ggtgttcgac ttttatattc gattctggtg gtttccaata atctcatgac	2760
atgcgtccgc ccgcatttat tgccagcgac ggccgggact tttccatccc tgggctgcta	2820
ggtcgggtac acgacccctcg ttttaccgcg aacgtaatgc tggaaagagc attgtgtcaa	2880
ggggggccgta gaagaagtc tccagcagag taaaatttcc tagggacggt aacggggcggt	2940
ggaaggagag agaagggaaag agacgtttct ggttccatta ctccacatTTT aagtttacc	3000
cgggagaatt ttacgttgcc agcagggtcg tccggatgtc aacggagcca gatatttcag	3060
agatggctac ctgattggac gaggacacca agacatttct acaaaaaaga tctgatctca	3120
tcgaccggct gcattaatga atcgccaaac ggcggggag aggccgttg cgtattggc	3180
gctttccgc ttctctcgatc actgactcg tgcgtcggtt cggtcgctg cggcgacgg	3240
tatcagctca ctcaaaggcg gtaatacggat tatccacaga atcagggat aacgcaggaa	3300
agaacatgtg agcaaaaaggc cagcaaaagg ccaggaacccg taaaaaggcc gcgttgctgg	3360
cggttttcca taggctccgc cccccctgacg agcatcacaa aaatcgacgc tcaagtccaga	3420
gggtggcgaaa cccgacacgaa ctataaaatgat accaggcggtt tccccctggaa agctccctcg	3480
tgcgtctcc tggcccgacc ctgccgttta ccggataacct gtccgccttt ctcccttgg	3540
gaagcgtggc gctttctcat agtcacgatc gtaggtatct cagttcggtg taggtcggtc	3600
gctccaagct gggctgtgtg cacgaacccc ccgttcagcc cgaccgtgc gccttatccg	3660
gttaactatcg tcttgagtc aacccggtaa gacacgactt atcgccactg gcagcagccaa	3720
ctggtaacag gattagcaga gcgaggatgtg taggcgggtgc tacagagttc ttgaagtgg	3780
ggcctaacta cggctacact agaaggacag tattttggat ctgcgtctg ctgaagccag	3840
ttaccttcgg aaaaagagtt ggtagctctt gatccggcaa acaaaccacc gctggtagcg	3900
gtggttttt tggcccgacc cagcagatcg cgcgcacaaa aaaaggatct caagaagatc	3960
ctttgtatctt ttctacgggg tctgacgctc agtggaaacga aaactcactt taagggattt	4020
tggtcatgat attatcaaaa aggtatctca cctagatcct tttaaattaa aatgaagtt	4080
ttaaatcaat cttaaagtata tatgagtaaa ctgggtctga cagttacca tgcttaatca	4140
gtgaggcacc tatctcagcg atotgtctat ttgcgttcatc catagttgcc tgactccccg	4200
tcgtgtatgat aactacgata cgggaggggct taccatctgg ccccaactgtc gcaatgatac	4260
cgccgagaccc acgctcaccg gctccagatt tatcagcaat aaaccagccaa gccggaaagg	4320
ccgagcgcag aagtggctt gcaactttat ccgcctccat ccagtctatt aattgttgcc	4380
ggaaagctag agtaagtagt tcgcccgttta atagtttgcg caacgttggtt gccattgcta	4440

-continued

caggcatcgt ggtgtcaegc	4500
gatcaaggcg agttacatga	4560
ctccgatcgt tgtcagaagt	4620
tgcataattc tcttactgtc	4680
caaccaagtc attctgagaa	4740
tacgggataa taccgcgeca	4800
cttcggggcg aaaactctca	4860
ctcgtgcacc caactgatct	4920
aaacaggaag gcaaaatgcc	4980
tcatactctt ccttttcaa	5040
attcgttatt gatttaaat	5100
tgatcattgt ttactaatta	5160
atttcagatg tctgaaaaag	5220
taattgcttc ataagcttca	5280
tggctacgccc tctttgcca	5340
tatcgtaact ctcttgcgtt	5400
aatctggagt taaaacttca	5460
cgaatttcgt atcagcaata	5520
acaactcgac cgccttgaca	5580
cttggtaat agagatgttt	5640
aaatcggagt tggaaaggcg	5700
ggacagtttt tgagttcttg	5760
tggcttccaa aggtatcgt	5820
atttgttatt ttcttcagac	5880
aagcaaccaa atgattcttt	5940
gagtcääat cttccctta	6000
tacggtcagt tgacgttgtt	6060
ttccttggc gattaaaggt	6120
gaaggataca gcaaagtgtg	6180
ctagagtagt atattggggc	6240
cttcaaagcg ccaagtggtt	6300
gtggagtatt taaaccatga	6360
tccgtattca taaacgatta	6420
tagtaacagt tatcatctt	6480
tgcgcactg ttggaaaggcg	6540
aggcacaaac gaacgtctca	6600
gaaaaacatc aaactcgaat	6660
gagataggtt gatcaggagc	6720
atactatata ggttacaaat	6780
tggcattctg acatcctttt	6840

-continued

```

acttgaactg aggaacagtc atgtctaagg ctaaaaactca cagttcgctg tggtcgacgg 6900
cgctattcag atcctcttct gagatgagtt ttgttgcgg cccaaaaata acttccaaat 6960
cttcagttat gtcaagcgat aggttaagaa tggtgatctt attatcattc tcataatgacg 7020
atctagtcgg caatgaactt tcttcgccc ttttttgac ttttataactg tctttcaacg 7080
ttatgttgtg ttgttaaatta ttctttgcga taatgatttt ttcaatattt gtattcctca 7140
aagtattgtg gtgaatgcct gtcattttctt cgggaatatg acttcgaaca ttttttaagg 7200
tattgttttca gaaaatgaac tgagtttcta 7230

```

```

<210> SEQ ID NO 26
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

```

```
<400> SEQUENCE: 26
```

```
gacgagatct tttttcaga ccatatgacc gg 32
```

```

<210> SEQ ID NO 27
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

```

```
<400> SEQUENCE: 27
```

```
gcgggattct tttctcagtt gatttgttg t 31
```

```

<210> SEQ ID NO 28
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

```

```
<400> SEQUENCE: 28
```

```
gcgggtcgac cacgacgaaac tgtgagttt agccttagac atgac 45
```

```

<210> SEQ ID NO 29
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

```

```
<400> SEQUENCE: 29
```

```
caggagcaaa gtcgtacga g 21
```

The invention claimed is:

1. A genetically engineered *Pichia* strain, which lacks a functional enzyme involved in production of high mannose structures, and comprises an exogenous nucleic acid encoding and expressing an exogenous enzyme for production of $\text{Man}_5\text{GlcNAc}_2$, wherein said enzyme involved in production of high mannose structures is α -1,6-mannosyltransferase encoded by the OCH1 gene, said exogenous enzyme for production of $\text{Man}_5\text{GlcNAc}_2$ is α -1,2-mannosidase or an enzymatically active fragment thereof and is targeted to the endoplasmic reticulum (ER), wherein said OCH1 gene is disrupted in said strain and the OCH1 gene disruption is the sole genetic disruption of genes coding for Golgi mannosyl

55 transferases acting in N-glycosylation of said strain, and wherein said strain produces $\text{Man}_5\text{GlcNAc}_2$ as a N-glycan structure or an intermediate N-glycan structure.

2. The strain of claim 1, wherein said exogenous enzyme is of a fungal origin or a mammalian origin.

3. The strain of claim 1, wherein the targeting of said exogenous enzyme to the ER is achieved by engineering said exogenous enzyme to include the ER retention signal as set forth in SEQ ID NO: 1.

4. A method for producing a glycoprotein with reduced hyperglycosylation in *Pichia*, comprising: (i) providing a *Pichia* strain, which lacks a functional enzyme involved in production of high mannose structures, and comprises an

exogenous nucleic acid encoding and expressing an exogenous enzyme for production of $\text{Man}_5\text{GlcNAc}_2$, wherein said enzyme involved in production of high mannose structures is α -1,6-mannosyltransferase encoded by the OCH1 gene, said exogenous enzyme for production of $\text{Man}_5\text{GlcNAc}_2$ is α -1, 5
2-mannosidase or an enzymatically active fragment thereof and is targeted to the ER, wherein said OCH1 gene is disrupted in said strain and the OCH1 gene disruption is the sole genetic disruption of genes coding for Golgi mannosyl transferases acting in N-glycosylation of said strain, and wherein 10
said strain produces $\text{Man}_5\text{GlcNAc}_2$ as a N-glycan structure or an intermediate N-glycan structure; and (ii) producing said glycoprotein in said *Pichia* strain.

5. The method of claim 4, wherein said exogenous enzyme is of a fungal origin or a mammalian origin. 15

6. The method of claim 4, wherein the targeting of said exogenous enzyme to the ER is achieved by engineering said exogenous enzyme to include the ER retention signal as set forth in SEQ ID NO: 1.

7. The strain of claim 1, wherein said α -1,2-mannosidase is 20
T. reesei α -1,2-mannosidase, and said strain produces $\text{Man}_5\text{GlcNAc}_2$ as a predominant N-glycan structure or a pre-dominant intermediate N-glycan structure.

8. The method of claim 4, wherein said α -1,2-mannosidase is *T. reesei* α -1,2-mannosidase, and said strain produces 25
 $\text{Man}_5\text{GlcNAc}_2$ as a predominant N-glycan structure or a pre-dominant intermediate N-glycan structure.

* * * *